

IPHC-2024-RAB025-05

INTERNATIONAL PACIFIC HALIBUT COMMISSION 5-YEAR PROGRAM OF INTEGRATED RESEARCH AND MONITORING (2022-26): UPDATES

PREPARED BY: IPHC SECRETARIAT (D. WILSON, J. PLANAS, I. STEWART, A. HICKS, B. HUTNICZAK, AND R. WEBSTER; 09 OCTOBER 2024)

PURPOSE

To provide the RAB with an annual opportunity to comment on the IPHC's 5-year Program of Integrated Research and Monitoring (2022-26) (the Plan).

BACKGROUND

Recalling that:

- the IPHC Secretariat conducts activities to address key issues identified by the Commission, its subsidiary bodies, the broader stakeholder community, and the IPHC Secretariat;
- b) the process of identifying, developing, and implementing the IPHC's science-based activities involves several steps that are circular and iterative in nature, but result in clear project activities and associated deliverables;
- c) the process includes developing and proposing projects based on direct input from the Commission, the experience of the IPHC Secretariat given its broad understanding of the resource and its associated fisheries, and concurrent consideration by relevant IPHC subsidiary bodies, and where deemed necessary, including by the Commission, additional external peer review;
- d) the IPHC Secretariat commenced implementation of the new Plan in 2022 and will keep the Plan under review on an ongoing basis.

Also recalling that an overarching goal of the IPHC 5-year Program of Integrated Research and Monitoring (2022-26) is to promote integration and synergies among the various research and monitoring activities of the IPHC Secretariat in order to improve knowledge of key inputs into the Pacific halibut stock assessment, and Management Strategy Evaluation (MSE) processes, thereby providing the best possible advice for management decision making processes.

The 1st iteration of the Plan was formally presented to the Commission at IM097 in November 2021 (IPHC-2021-IM097-12) for general awareness of the documents ongoing development. At the 98th Session of the IPHC Annual Meeting (AM098) in January 2022, the Commission requested a number of amendments which were subsequently incorporated.

In 2023 and 2024, the plan went through two cycles of review and improvement with the SRB, with amendments being suggested and incorporated accordingly.

DISCUSSION

The RAB should note that:

- a) the intention is to ensure that the new integrated plan is kept as a 'living plan', and is reviewed and updated annually based on the resources available to undertake the work of the Commission (e.g. internal and external fiscal resources, collaborations, internal expertise);
- b) the plan focuses on core responsibilities of the Commission; and any redirection provided by the Commission;



IPHC-2024-RAB025-05

c) each year the SRB may choose to recommend modifications to the current Plan, and that any modifications subsequently made would be documented both in the Plan itself, and through reporting back to the SRB and then the Commission.

Updates: The Secretariat is currently in the process of updating the Plan to meet the request of the SRB at it's 24th Session, as per the below text:

International Pacific Halibut Commission 5-year program of integrated research and monitoring (2022-26)

SRB024—Req.01 (para. 14) The SRB **REQUESTED** that the IPHC 5-year Program of Integrated Research and Monitoring be revised by SRB026 to reflect changing priorities in light of major progress on biological research and ongoing monitoring challenges.

RECOMMENDATION

That the RAB:

1) **NOTE** paper IPHC-2024-RAB025-05 that provides the latest iteration of the IPHC 5-year program of Integrated Research and Monitoring (2022-26).

APPENDICES

Appendix A: Updated: IPHC 5-year program of Integrated Research and Monitoring (2022-26)

INTERNATIONAL PACIFIC HALIBUT COMMISSION 5-YEAR PROGRAM OF INTEGRATED RESEARCH AND MONITORING

(2022 - 2026)

INTERNATIONAL PACIFIC



Commissioners

Canada United States of America

Paul Ryall Jon Kurland

Neil Davis Robert Alverson

Peter DeGreef Richard Yamada

Executive Director

David T. Wilson, Ph.D.

BIBLIOGRAPHIC ENTRY

IPHC 2023. International Pacific Halibut Commission 5-Year program of integrated research and monitoring (2022-26). Seattle, WA, U.S.A. *IPHC*–2023-5YPIRM, 58 pp.





The designations employed and the presentation of material in this publication and its lists do not imply the expression of any opinion whatsoever on the part of the International Pacific Halibut Commission (IPHC) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

This work is protected by copyright. Fair use of this material for scholarship, research, news reporting, criticism or commentary is permitted. Selected passages, tables or diagrams may be reproduced for such purposes provided acknowledgment of the source is included. Major extracts or the entire document may not be reproduced by any process without the written permission of the Executive Director, IPHC.

The IPHC has exercised due care and skill in the preparation and compilation of the information and data set out in this publication. Notwithstanding, the IPHC, its employees and advisers, assert all rights and immunities, and disclaim all liability, including liability for negligence, for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying upon any of the information or data set out in this publication, to the maximum extent permitted by law including the International Organizations Immunities Act.

Contact details:

International Pacific Halibut Commission 2320 W. Commodore Way, Suite 300

Seattle, WA, 98199-1287, U.S.A.

Phone: +1 206 634 1838 Fax: +1 206 632 2983

Email: secretariat@iphc.int
Website: http://www.iphc.int/

ACRONYMS

AM Annual Meeting
CB Conference Board
DMR Discard Mortality Rate

FAC Finance and Administration Committee FISS Fishery-Independent Setline Survey

FSC First Nations Food, Social, and Ceremonial [fishery]

IM Interim Meeting

IPHC International Pacific Halibut Commission
MSAB Management Strategy Advisory Board
MSE Management Strategy Evaluation

OM Operating Model

PAB Processor Advisory Board PDO Pacific Decadal Oscillation

PHMEIA Pacific halibut multiregional economic impact assessment [model]

QAQC Quality assurance/quality control

RAB Research Advisory Board

SHARC Subsistence Halibut Registration Certificates

SRB Scientific Review Board

TCEY Total Constant Exploitation Yield

U.S.A. United States of America

WM Work Meeting

DEFINITIONS

A set of working definitions are provided in the IPHC Glossary of Terms and abbreviations: https://iphc.int/the-commission/glossary-of-terms-and-abbreviations



TABLE OF CONTENTS

1. Introduction	
2. Objectives	8
3. Strategy	11
4. Measures of Success	
4.1 Delivery of specified products	12
4.2 Communication	13
4.3 External research funding	13
4.4 Peer-reviewed journal publication	13
5. Core focal areas – Background	13
5.1 Research	14
5.1.1 Stock Assessment	14
5.1.2 Management Strategy Evaluation (MSE)	
5.1.3 Biology and Ecology	16
5.2 Monitoring	
5.2.1 Fishery-dependent data	
5.2.1.1 Directed commercial fisheries data	
5.2.1.2 Non-directed commercial discard mortality data	18
5.2.1.3 Subsistence fisheries data	18
5.2.1.4 Recreational fisheries data	18
5.2.2 Fishery-independent data	19
5.2.2.1 Fishery-independent setline survey (FISS)	19
5.2.2.2 Fishery-independent Trawl Survey (FITS)	20
5.2.2.3 Norton Sound trawl survey	21
5.2.3 Age composition data (both fishery-dependent and fishery-independe	nt)21
5.3 Management-supporting information	21
6. Core focal areas – Planned and opportunistic activities (2022-2026)	
6.1 Research	
6.1.1 Stock Assessment	
6.1.1.1 Stock Assessment data collection and processing	
6.1.1.2 Stock Assessment technical development	
6.1.1.3 Stock Assessment biological inputs	
6.1.1.4 Stock Assessment fishery yield	26



6.1.2	Management Strategy Evaluation	27	
6.1.2.1	MSE Biological and population parameterization	27	
6.1.2.2	MSE technical development	28	
6.1.2.3	MSE Program of Work for 2021–2023	29	
6.1.2.4	Potential Future MSE projects	29	
6.1.3	Biology and Ecology	29	
6.1.3.1	Migration and Population Dynamics	30	
6.1.3.2 Reproduction		30	
6.1.3.3	Growth	31	
6.1.3.4	6.1.3.4 Mortality and Survival Assessment		
6.1.3.5	Fishing Technology	31	
6.2 Mo	onitoring	31	
6.2.1	Fishery-dependent data	32	
6.2.1.1	Directed commercial fisheries data	32	
	Annually review the spatial distribution of sampling effort among ports, data collections rates, and quality assurance/quality control (QAQC) processes, including in-season regactivities	view of port	
6.2.1.3			
6.2.1.4	Recreational fisheries data	32	
6.2.2	Fishery-independent data	32	
6.2.2.1	Fishery-independent setline survey (FISS)	32	
6.2.2.2	Fishery-independent Trawl Survey (FITS)	33	
6.2.3	Age composition data (both fishery-dependent and fishery-independent)	33	
6.3 Po	tential of integrating human dynamics into management decision-making	33	
7. Amend	lment	34	
3. Refere	nces	35	

EXECUTIVE SUMMARY

An overarching goal of the *IPHC 5-Year Program of Integrated Research and Monitoring (2022-26)* is to promote integration and synergies among the various research and support activities of the IPHC Secretariat in order to improve our knowledge of key inputs into the Pacific halibut stock assessment and Management Strategy Evaluation (MSE) processes, and to provide the best possible advice for management decision-making processes.

Along with the implementation of the short- and medium-term activities contemplated in this *IPHC 5-Year Program of Integrated Research and Monitoring (2022-26)*, and in pursuit of the overarching objective, the IPHC Secretariat will also aim to:

- 1) undertake cutting-edge research programs in fisheries research in support of Pacific halibut fisheries management;
- 2) undertake groundbreaking methodological research;
- 3) undertake applied research;
- 4) establish new collaborative agreements and interactions with research agencies and academic institutions;
- 5) promote the international involvement of the IPHC by continued and new participation in international scientific organizations and by leading international science and research collaborations;
- 6) effectively communicate IPHC research outcomes;
- 7) incorporate talented students and early researchers in research activities contemplated.

The research and monitoring activities conducted by the IPHC Secretariat are directed towards fulfilling the following four (4) objectives within areas of data collection, biological and ecological research, stock assessment, and Management Strategy Evaluation (MSE). In addition, the IPHC responds to Commission requests for additional inputs to management and policy development which are classified under management support.

The Secretariat's success in implementing the *IPHC 5-Year Program of Integrated Research and Monitoring* (2022-26) will be measured according to the following criteria relevant to the stock assessment, the MSE and for all inputs to IPHC management:

- 1) Timeliness was the research conducted, analyzed, published, and provided to the Commission at the appropriate points to be included in annual management decisions?
- 2) Accessibility was the research published and presented in such a way that it was available to other scientists, stakeholders, and decision-makers?
- 3) Relevance did the research improve the perceived accuracy of the stock assessment, MSE, or decisions made by the Commission?
- 4) Impact did the research allow for more precision or a better estimate of the uncertainty associated with information for use in management?
- 5) Reliability has the research resulted in more consistent information provided to the Commission for decision-making?

1. Introduction

The International Pacific Halibut Commission (IPHC) is a public international organization so designated via Presidential Executive Order 11059 and established by a Convention between Canada and the United States of America. The IPHC Convention was signed on 2 March 1923, ratified on 21 July 1924, and came into effect on 21 October 1924 upon exchange. The Convention has been revised several times since, to extend the Commission's authority and meet new conditions in the fishery. The most recent change occurred in 1979 and involved an amendment to the 1953 Halibut Convention. The 1979 amendment, termed a "protocol", was precipitated in 1976 by Canada and the United States of America extending their jurisdiction over fisheries resources to 200 miles. The 1979 Protocol along with the U.S. legislation that gave effect to the Protocol (Northern Pacific Halibut Act of 1982) has affected the way the fisheries are conducted, and redefined the role of IPHC in the management of the fishery. Canada does not require specific enabling legislation to implement the protocol.

The basic texts of the Commission are available on the IPHC website: https://www.iphc.int/the-commission, and prescribe the mission of the organization as:

".... to develop the stocks of [Pacific] halibut in the Convention waters to those levels which will permit the optimum yield from the fishery and to maintain the stocks at those levels." IPHC Convention, Article I, sub-article I, para. 2). The IPHC Convention Area is detailed in Fig. 1.

The IPHC Secretariat, formed in support the Commission's activities, is based in Seattle, WA, U.S.A. As its shared vision, the IPHC Secretariat aims to deliver positive economic, environmental, and social outcomes for the Pacific halibut resource for Canada and the U.S.A. through the application of rigorous science, innovation, and the implementation of international best practice.




Figure 1. Map of the IPHC Convention Area (map insert) and IPHC Regulatory Areas.

2. Objectives

The IPHC has a long-standing history (since 1923) of collecting data, undertaking research, and stock assessment, devoted to describing and understanding the Pacific halibut (*Hippoglossus stenolepis*) stock and the fisheries that interact with it.

The IPHC Secretariat conducts activities to address key issues identified by the Commission, its subsidiary bodies, the broader stakeholder community, and of course, the IPHC Secretariat itself. The process of identifying, developing, and implementing our science-based activities involves several steps that are circular in nature, but result in clear research activities and associated deliverables. The process includes developing and proposing projects based on direct input from the Commission, the experience of the IPHC Secretariat given our broad understanding of the resource and its associated fisheries, and concurrent consideration by relevant IPHC subsidiary bodies, and where deemed necessary, additional external peer review.

Over the last five years (2017-2021), the research conducted by the IPHC Secretariat has been guided by a 5-Year Biological and Ecosystem Science Research Plan (IPHC-2019-BESRP-5YP) that aimed at improving knowledge on the biology of Pacific halibut in order to improve the accuracy of the stock assessment and in the management strategy evaluation (MSE) process. The IPHC-2019-BESRP-5YP contemplated research activities in five focal areas, namely Migration and Distribution, Reproduction, Growth and Physiological Condition, Discard Mortality Rates and Survival, and Genetics and Genomics. Research activities were highly integrated with the needs of stock assessment and MSE by their careful alignment with biological uncertainties and parameters, and the resulting prioritization (Appendix I). The outcomes of the IPHC-2019-BESRP-5YP have provided key inputs into stock assessment and the MSE process and, importantly, have provided foundational information for the successful pursuit of continuing and novel objectives within the new 5-Year Program of Integrated Research and Monitoring (2022-2026) (5YPIRM) (Appendix I).

The 2nd Performance Review of the IPHC (<u>IPHC-2019-PRIPHC02-R</u>), carried out over the course of 2019, also provided a range of recommendations to the Commission on ways in which it could continue to improve on the quality of scientific advice being provided to the Commission. There were nine (9) specific recommendations as provided below:

Science: Status of living marine resources

PRIPHC02–Rec.03 (para. 44) The PRIPHC02 **RECOMMENDED** that opportunities to engage with western Pacific halibut science and management agencies be sought, to strengthen science links and data exchange. Specifically, consider options to investigate pan-Pacific stock structure and migration of Pacific halibut.

PRIPHC02–Rec.04 (para. 45) The PRIPHC02 RECOMMENDED that:

- *a) further efforts be made to lead and collaborate on research to assess the ecosystem impacts of Pacific halibut fisheries on incidentally caught species (retained and/or discarded);*
- b) where feasible, this research be incorporated within the IPHC's 5-Year Research Plan (https://www.iphc.int/uploads/pdf/besrp/2019/iphc-2019-besrp-5yp.pdf);
- c) findings from the IPHC Secretariat research and that of the Contracting Parties be readily accessible via the IPHC website.

Science: Quality and provision of scientific advice

PRIPHC02–Rec.05 (para. 63) The PRIPHC02 **RECOMMENDED** that simplified materials be developed for RAB and especially MSAB use, including training/induction materials.

PRIPHC02—Rec.06 (para. 64) The PRIPHC02 **RECOMMENDED** that consideration be given to amending the Rules of Procedure to include appropriate fixed terms of service to ensure SRB peer review remains independent and fresh; a fixed term of three years seems appropriate, with no more than one renewal.

PRIPHC02–Rec.07 (para. 65) The PRIPHC02 **RECOMMENDED** that the peer review process be strengthened through expanded subject specific independent reviews including data quality and standards, the FISS, MSE, and biological/ecological research; as well as conversion of "grey literature" to primary literature publications. The latter considered important to ongoing information outreach efforts given the cutting-edge nature of the Commission's scientific work.

PRIPHC02–Rec.08 (para. 66) The PRIPHC02 **RECOMMENDED** that the IPHC Secretariat develop options for simple graphical summaries (i.e. phase plot equivalents) of fishing intensity and spawning stock biomass for provision to the Commission.

Conservation and Management: Data collection and sharing

PRIPHC02–Rec.09 (para. 73) The PRIPHC02 **RECOMMENDED** that observer coverage be adjusted to be commensurate with the level of fishing intensity in each IPHC Regulatory Area.

Conservation and Management: Consistency between scientific advice and fishery Regulations adopted

PRIPHC02—Rec.10 (para. 82) The PRIPHC02 **RECOMMENDED** that the development of MSE to underpin multi-year (strategic) decision-making be continued, and as multi-year decision making is implemented, current Secretariat capacity usage for annual stock assessments should be refocused on research to investigate MSE operating model development (including consideration of biological and fishery uncertainties) for future MSE iterations and regularised multi-year stock assessments.

PRIPHC02—Rec.11 (para. 83) The PRIPHC02 **RECOMMENDED** that ongoing work on the MSE process be prioritised to ensure there is a management framework/procedure with minimal room for ambiguous interpretation, and robust pre-agreed mortality limit setting frameworks.

The work outlined in this document builds on the previous a 5-Year Biological and Ecosystem Science Research Plan (IPHC-2019-BESRP-5YP), closing completed projects, extending efforts where needed, and adding new avenues in response to new information. Appendix I provides a detailed summary of the previous plan and the status of the work specifically undertaken. Key highlights relevant to the stock assessment and MSE include:

- Completion of the genetic assay for determining sex from tissue samples, processing of commercial fishery samples collected during 2017-2020, inclusion of this information in the 2019 and subsequent stock assessments, and transfer of this effort from research to ongoing monitoring.
- Incremental progress toward population-level sampling and analysis of maturity and fecundity.
- Continued development of the understanding of physiological and environmental mechanisms determining growth for future field application.
- Published estimates of discard mortality rates for use in data processing and management accounting.
- Collection of genetic samples and genome sequencing to provide a basis for ongoing evaluation of stock structure at population-level and finer scales.

All previously described research areas continue to represent critical areas of uncertainty in the stock assessment and thus are closely linked to management performance. The previous 5-year plan was successful in either providing direct new information to the stock assessment or building the foundation for the collection/analysis

of such information in this updated plan. As noted below, some new priorities have emerged, and others have evolved based on the work completed to date. The incorporation of research objectives in the 5YPIRM that address climate change as a factor influencing Pacific halibut biology and ecology as well as fishery performance and dynamics constitutes a timely and relevant contribution towards advancing IPHC-led research to the forefront of fisheries science.

An **overarching goal** of the *IPHC 5-Year Program of integrated research and monitoring (2022-26)* is therefore to promote integration and synergies among the various research and support activities of the IPHC Secretariat in order to improve our knowledge of key inputs into the Pacific halibut stock assessment and MSE processes, in order to provide the best possible advice for management decision-making processes.

Along with the implementation of the short- and medium-term activities contemplated in this *IPHC 5-Year Program of Integrated Research and monitoring (2022-26)*, and in pursuit of the overarching objective, the IPHC Secretariat will also aim to:

- 1) undertake cutting-edge research programs in fisheries research in support of fisheries management of Pacific halibut;
- 2) undertake groundbreaking methodological research;
- 3) undertake applied research;
- 4) establish new collaborative agreements and interactions with research agencies and academic institutions;
- 5) promote the international involvement of the IPHC by continued and new participation in international scientific organizations and by leading international science and research collaborations.
- 6) effectively communicate IPHC research outcomes
- 7) incorporate talented students and early researchers in research activities contemplated.

The research and monitoring activities conducted by the IPHC Secretariat are directed towards fulfilling the following four (4) **objectives** within areas of data collection, biological and ecological research, stock assessment, and MSE. In addition, the IPHC responds to Commission requests for additional inputs to management and policy development which are classified under management support. The overall aim is to provide a program of integrated research and monitoring (Fig 2):

Research

- 1) Stock assessment: apply the resulting knowledge to improve the accuracy and reliability of the current stock assessment and the characterization of uncertainty in the resultant stock management advice provided to the Commission;
- 2) Management Strategy Evaluation (MSE): to develop an accurate, reliable, and informative MSE process to appropriately characterize uncertainty and provide for the robust evaluation of the consequences of alternative management options, known as harvest strategies, using defined conservation and fishery objectives;
- 3) <u>Biology and Ecology</u>: identify and assess critical knowledge gaps in the biology and ecology of Pacific halibut within its known range, including the influence of environmental conditions on population and fishery dynamics;



Monitoring

4) **Monitoring**: collect representative fishery dependent and fishery-independent data on the distribution, abundance, biology, and demographics of Pacific halibut through ongoing monitoring activities;

Integrated management support

5) <u>Additional management-supporting inputs</u>: respond to Commission requests for any additional information supporting management and policy development.




Figure 2. Core areas of the IPHC's program of integrated research and monitoring providing management support.

3. Strategy

The IPHC Secretariat has five (5) enduring strategic goals in executing our mission, including our overarching goal and associated science and research objectives, as articulated in our Strategic Plan (IPHC Strategic Plan (2019-23)): 1) To operate in accordance with international best practice; 2) Be a world leader in scientific excellence and science-based decision making; 3) To foster collaboration (within Contracting Parties and internationally) to enhance our science and management advice; 4) Create a vibrant IPHC culture; and 5) Set the

standard for fisheries commissions globally.

Although priorities and tasking will change over time in response to events and developments, the Strategic Plan provides a framework to standardise our approach when revising or setting new priorities and tasking. The Strategic goals as they apply to the science and research activities of the IPHC Secretariat, will be operationalised through a multi-year tactical activity matrix at the organisational and management unit (Branch) level (Fig. 3). The tactical activity matrix is described in the sections below and has been developed based on the core needs of the Commission, in developing and implementing robust, scientifically-based management decisions on an annual, and multi-year level. Relevant IPHC subsidiary bodies will be involved in project development and ongoing review.




Figure 3. IPHC Secretariat organisation chart (2023).

4. Measures of Success

The Secretariat's success in implementing the *IPHC 5-Year Program of Integrated Research and Monitoring* (2022-26) will be measured according to the following criteria relevant to the stock assessment, the MSE and for all inputs to IPHC management:

- 1) Timeliness was the research conducted, analyzed, published, and provided to the Commission at the appropriate points to be included in annual management decisions?
- 2) Accessibility was the research published and presented in such a way that it was available to other scientists, stakeholders, and decision-makers?
- 3) Relevance did the research improve the perceived accuracy of the stock assessment, MSE or decisions made by the commission?
- 4) Impact did the research allow for more precision or a better estimate of the uncertainty associated with information for use in management?
- 5) Reliability has research resulted in more consistent information provided to the Commission for decision-making.

4.1 Delivery of specified products

Each project line item will contain specific deliverables that constitute useful inputs into the stock assessment and the management strategy evaluation process, as well as support their implementation in the decision-making

process at the level of the Commission.

4.2 Communication

The IPHC Secretariat will disseminate information about the activities contemplated in the IPHC 5-Year Program of Integrated Research and Monitoring (2022-2026) and the resulting products to Contracting Parties, stakeholders, the scientific community, and the general public through a variety of channels:

- 1) IPHC website (www.iphc.int);
- 2) Formal documentation provided for IPHC meetings (Interim and Annual Meetings, Subsidiary Body meetings, etc.);
- 3) Presentations at national and international scientific conferences;
- 4) Published reports and peer-reviewed publications (section 4.4);
- 5) Outreach events;
- 6) Social media outlets (e.g. Facebook, Twitter, LinkedIn, etc.);
- 7) Informal presentations and interactions with partners, stakeholders, and decision-makers at varied times and venues when needed.

4.3 External research funding

The Secretariat has set a funding goal of at least 20% of the funds for this program to be sourced from external funding bodies on an annual basis. Continuing the successful funding-recruitment strategy adopted during the previous 5-yr research plan (IPHC–2019–BESRP-5YP) (Appendix I), the Secretariat will identify and select external funding opportunities that are timely and that aim at addressing key research objectives (as outlined in Appendix II and summarized in Appendix V) that have important implications for stock assessment and the MSE process. The IPHC Secretariat has the necessary expertise to propose novel and important research questions to funding agencies and to recruit external collaborators from research agencies and universities as deemed necessary. The IPHC Secretariat will continue to capitalize on the strong analytical contributions of quantitative scientists to the development of biological research questions within the framework of research projects funded by external as well as internal funding sources.

4.4 Peer-reviewed journal publication

Publication of research outcomes in peer-reviewed journals will be clearly documented and monitored as a measure of success. This may include single publications at the completion of a particular project, or a series of publications throughout the project as well as at its completion. Each sub-project shall be published in a timely manner and shall be submitted no later than 12 months after the end of the research. In the sections that follow, the expected publications from each research stream and cross-stream are defined.

5. Core focal areas – Background

The goals of the main activities of the 5-Year program of integrated research and monitoring (2022-26) are integrated across the organisation, involving 1) monitoring (fisheries-dependent and –independent data collection), and 2) research (biological, ecological), modelling (FISS and stock assessment), and MSE, as outlined in the following sub-sections. These components are closely linked to one another, and all feed into management decision-making (Fig. 4). Additionally, management-supporting information constitute a range of additional decision-making drivers within and beyond IPHC's current research and monitoring programs. The current program builds on the outcomes and experiences of the Commission arising from the implementation of the 2017-21 5-Year Biological and Ecosystem Science Research Plan (IPHC-2019-BESRP-5YP), and which is summarized in Appendix I.




Figure 4. Flow of information from basic biological understanding of the Pacific halibut resource, through IPHC research components (monitoring, biological and ecological research, stock assessment, and MSE) to management decision-making. Management-supporting information (grey) constitute a range of additional decision-making drivers within and beyond IPHC's current research and monitoring programs. Arrows indicate the strength (size of the arrow) and direction of information exchange. Also identified (in black) are the external links from funding and scientific publications which supplement the IPHC's internal process.

5.1 Research

5.1.1 Stock Assessment

Focal Area Objective	To improve accuracy and reliability of the current stock assessment and the characterization of uncertainty in the resultant stock management advice provided to the Commission.	
IPHC Website portal	https://www.iphc.int/management/science-and-research/stock-assessment	

The IPHC conducts an annual stock assessment, using data from the fishery-independent setline survey (FISS), the commercial Pacific halibut and other fisheries, as well biological information from its research program. The assessment includes the Pacific halibut resource in the IPHC Convention Area, covering the Exclusive Economic Zones of Canada and the United States of America. Data sources are updated each year to reflect the most recent scientific information available for use in management decision-making.

The 2021 stock assessment relied on an ensemble of four population dynamics models to estimate the probability distributions describing the current stock size, trend, and demographics. The ensemble is designed to capture both uncertainty related to the data and stock dynamics (due to estimation) as well as uncertainty related to our understanding of the way in which the Pacific halibut stock functions and is best approximated by a statistical model (structural uncertainty).

Stock assessment results are used as inputs for harvest strategy calculations, including mortality projection tables for the upcoming year that reflect the IPHC's harvest strategy policy and other considerations, as well as the

harvest decision table which provides a direct tool for the management process. The harvest decision table uses the probability distributions from short-term (three year) assessment projections to evaluate the trade-offs between alternative levels of potential yield (catch) and the associated risks to the stock and fishery.

The stock assessment research priorities have been subdivided into four categories:

- 1) Assessment data collection and processing;
- 2) technical development;
- 3) biological inputs; and
- 4) fishery yield.

It is important to note that ongoing monitoring, including the annual FISS and directed commercial landings sampling programs is not considered research and is therefore not included in this research priority list despite the critical importance of these collections. These are described in the sections below.

5.1.2 Management Strategy Evaluation (MSE)

Focal Area Objective	To develop an accurate, reliable, and informative MSE process to appropriately characterize uncertainty and provide for the robust evaluation of the consequences of alternative management options, known as harvest strategies, using defined conservation and fishery objectives.	
IPHC Website portal https://www.iphc.int/management/science-and-research/management-strategy evaluation		

Management Strategy Evaluation (MSE) is a process to evaluate the consequences of alternative management options, known as harvest strategies. MSE uses a simulation tool to determine how alternative harvest strategies perform given a set of pre-defined fishery and conservation objectives, taking into account the uncertainties in the system and how likely candidate harvest strategies are to achieve the chosen management objectives.

MSE is a simulation technique based on modelling each part of a management cycle. The MSE uses an operating model to simulate the entire population and all fisheries, factoring in management decisions, the monitoring program, the estimation model, and potential ecosystem effects using a closed-loop simulation.

Undertaking an MSE has the advantage of being able to reveal the trade-offs among a range of possible management decisions. Specifically, to provide the information on which to base a rational decision, given harvest strategies, preferences, and attitudes to risk. The MSE is an essential part of the process of developing, evaluating and agreeing to a harvest strategy.

The MSE process involves:

- Defining fishery and conservation objectives with the involvement of stakeholders and managers;
- Identifying harvest strategies (a.k.a. management procedures) to evaluate;
- Simulating a Pacific halibut population using those harvest strategies;
- Evaluating and presenting the results in a way that examines trade-offs between objectives;
- Applying a chosen harvest strategy for the management of Pacific halibut;
- Repeating this process in the future in case of changes in objectives, assumptions, or expectations.

There are many tasks that would continue to improve the MSE framework and the presentation of future results to the Commission. The tasks can be divided into five general categories, which are common to MSE in general:

- 1. **Objectives**: The goals and objectives that are used in the evaluation.
- 2. **Management Procedures (MPs)**: Specific, well-defined management procedures that can be coded in the MSE framework to produce simulated Total Constant Exploitation Yields (TCEY) for each IPHC Regulatory Area.
- 3. **Framework**: The specifications and computer code for the closed-loop simulations including the operating model and how it interacts with the MP.
- 4. **Evaluation**: The performance metrics and presentation of results. This includes how the performance metrics are evaluated (e.g. tables, figures, and rankings), presented to the Commission and its subsidiary bodies, and disseminated for outreach.
- 5. **Application**: Specifications of how an MP may be applied in practice and re-evaluated in the future, including responses to exceptional circumstances.

All these categories provide inputs and outputs of the MSE process, but the Framework category benefits most from the integration of biological and ecosystem research because the operating model, the simulation of the monitoring program, the estimation model, and potential ecosystem effects are determined from this knowledge.

Outcomes of the MSE process will not only inform the Commission on trade-offs between harvest strategies and assist in choosing an optimal strategy for management of the Pacific halibut resource but will inform the prioritization of research activities related to fisheries monitoring, biological and ecological research, stock assessment, and fishery socioeconomics.

5.1.3 Biology and Ecology

Focal Area Objective To identify and assess critical knowledge gaps in the biology and ecology halibut within its known range, including the influence of environmental on population and fishery dynamics.	
IPHC Website portal	https://www.iphc.int/management/science-and-research/biological-and-ecosystem-science-research-program-bandesrp

Since its inception, the IPHC has had a long history of research activities devoted to describe and understand the biology of the Pacific halibut. At present, the main objectives of the Biological and Ecosystem Science Research Program at IPHC are to: 1) identify and assess critical knowledge gaps in the biology of the Pacific halibut; 2) understand the influence of environmental conditions in the biology of the Pacific halibut and its fishery; and 3) apply the resulting knowledge to reduce uncertainty in current stock assessment models.

The primary biological research activities at the IPHC that follow Commission objectives and that are selected for their important management implications are identified and described in the proposed IPHC 5-Year Program of Integrated Research and Monitoring (2022-2026). An overarching goal of the 5-Year Program of Integrated Research and Monitoring (2022-2026) is to promote integration and synergies among the various research activities led by the IPHC to improve our knowledge of key biological inputs that feed into the stock assessment and MSE process. The goals of the main research activities of the 5-Year Program of Integrated Research and Monitoring (2022-2026) are therefore aligned and integrated with the IPHC stock assessment and MSE processes. The IPHC Secretariat conducts research activities to address key biological issues based on the IPHC Secretariat's own input as well as input from the IPHC Commissioners, stakeholders and particularly from specific subsidiary

bodies to the IPHC, including the Scientific Review Board (SRB) and the Research Advisory Board (RAB).

The biological research activities contemplated in the 5-Year Program of Integrated Research and Monitoring (2022-2026) and their specific aims are detailed in Section 6. Overall, the biological research activities at the IPHC aim to provide information on 1) factors that influence the biomass of the Pacific halibut population (e.g. distribution and movement of fish among IPHC Regulatory Areas, growth patterns and environmental influences on growth in larval, juvenile and adult fish, drivers of changes in size-at-age); 2) the spawning (female) population (e.g. reproductive maturity, skipped spawning, reproductive migrations); and 3) resulting changes in population dynamics. Furthermore, the research activities of IPHC also aim to provide information on the survival of regulatory-discarded Pacific halibut in the directed fisheries with the objective to refine current estimates of discard mortality rates and develop best handling practices, and reduce whale depredation and Pacific halibut bycatch through gear modifications and through a better understanding of behavioral and physiological responses of Pacific halibut to fishing gear. The proposed timeline of activities and of staffing and funding indicators are provided in Appendix VI and Appendix VII, respectively.

5.2 Monitoring

Focal Area Objective	To collect fishery-dependent and fishery-independent data on the distribution, abundance, and demographics of Pacific halibut, as well as other key biological data, through ongoing monitoring activities.	
Fishery-dependent data:		
IPHC Website portal	• https://www.iphc.int/datatest/commercial-fisheries	
	• https://www.iphc.int/data/datatest/pacific-halibut-recreational-fisheries-data	
	• https://www.iphc.int/datatest/subsistence-fisheries	
	• https://www.iphc.int/data/time-series-datasets	
	Fishery-independent data:	
	 https://www.iphc.int/management/science-and-research/fishery- independent-setline-survey-fiss 	
	• https://www.iphc.int/data/datatest/fishery-independent-setline-survey-fiss	
	• https://www.iphc.int/datatest/data/water-column-profiler-data	

5.2.1 Fishery-dependent data

The IPHC estimates all Pacific halibut removals taken in the IPHC Convention Area and uses this information in its yearly stock assessment and other analyses. The data are compiled by the IPHC Secretariat and include data from Federal and State agencies of each Contracting Party. Specific activities in this area are described below.

5.2.1.1 Directed commercial fisheries data

The IPHC Secretariat collects logbooks, otoliths, tissue samples, and associated sex-length-weight data from directed commercial landings coastwide (Fig. 5). A sampling rate is determined for each port by IPHC Regulatory Area. The applicable rate is calculated from the current year's mortality limits and estimated percentages of weight of fish landed, and estimated percentages of weight sampled in that port to allow for collection of the target number of biological samples by IPHC Regulatory Area. An example of the data collected and the methods used are provided in the annually updated directed commercial sampling manual (e.g. IPHC Directed Commercial Landings Sampling Manual 2022). Directed commercial fishery landings are recorded by the Federal and State agencies of each Contracting Party and summarized each year by the IPHC. Discard mortality for the directed

commercial fishery is currently estimated using a combination of research survey (U.S.A.) and observer data (Canada).

5.2.1.2 Non-directed commercial discard mortality data

The IPHC accounts for non-directed commercial discard mortality by IPHC Regulatory Area and sector. Non-directed commercial discard mortality estimates are provided by State and Federal agencies of each Contracting Party and compiled annually for use in the stock assessment and other analyses.

Non-directed commercial discard mortality of Pacific halibut is estimated because not all fisheries have 100% monitoring and not all Pacific halibut that are discarded are assumed to die. The IPHC relies upon information supplied by observer programs run by Contracting Party agencies for non-directed commercial discard mortality estimates in most fisheries. Non-IPHC research survey information or other sources are used to generate estimates of non-directed commercial discard mortality in the few cases where fishery observations are unavailable. Non-directed fisheries off Canada British Columbia are monitored and discard mortality information is provided to IPHC by DFO. NOAA Fisheries operates observer programs off the USA West Coast and Alaska, which monitor the major groundfish fisheries. Data collected by those programs are used to estimate non-directed commercial discard mortality.

5.2.1.3 Subsistence fisheries data

Subsistence fisheries are non-commercial, customary, and traditional use of Pacific halibut for direct personal, family, or community consumption or sharing as food, or customary trade. The primary subsistence fisheries are the treaty Indian Ceremonial and Subsistence fishery in IPHC Regulatory Area 2A off northwest Washington State (USA), the First Nations Food, Social, and Ceremonial (FSC) fishery in British Columbia (Canada), and the subsistence fishery by rural residents and federally recognized native tribes in Alaska (USA) documented via Subsistence Halibut Registration Certificates (SHARC). Subsistence fishery removals of Pacific halibut, including estimated subsistence discard mortality, are provided by State and Federal agencies of each Contracting Party, estimated, and compiled annually for use in the stock assessment and other analysis.

5.2.1.4 Recreational fisheries data

Recreational removals of Pacific halibut, including estimated recreational discard mortality, are provided by National/State agencies of each Contracting Party, estimated, and compiled annually for use in the stock assessment and other analysis.

IPHC 5-Year program of integrated research and monitoring (2022-26)




Figure 5. Ports where the IPHC has sampled directed commercial landings throughout the fishing period in recent years (note: ports sampled may change from year-to-year for operational reasons).

5.2.2 Fishery-independent data

Data collection and monitoring activities aimed at providing a standardised time-series of biological and ecological data that is independent of the fishing fleet.

5.2.2.1 Fishery-independent setline survey (FISS)

The IPHC Fishery-Independent Setline Survey (FISS) provides catch-rate information and biological data on Pacific halibut that are independent of the fishery. These data, collected using standardized methods, bait, and gear, are used to estimate the primary index of population abundance used in the stock assessment. The FISS is restricted to the summer months but encompasses the commercial fishing grounds in the Pacific halibut fishery, and almost all known Pacific halibut habitat in Convention waters outside the Bering Sea. The standard FISS grid totals 1,890 stations (Fig. 6). Biological data collected on the FISS (e.g. the length, weight, age, and sex of Pacific halibut) are used to monitor changes in biomass, growth, and mortality. In addition, records of non-target species caught during FISS operations provide insight into bait competition, and serve as an index of abundance over time, making them valuable to the potential management and avoidance of non-target species. Environmental data are also collected including water column temperature, salinity, dissolved oxygen, pH, and chlorophyll concentration to help identify the conditions in which the fish were caught, and these data can serve as co-variates in space-time modeling used in the stock assessment. An example of the data collected and the methods used are provided in the annually updated FISS sampling manual (e.g. IPHC FISS Sampling Manual 2022).




Figure 6. IPHC Fishery-Independent Setline Survey (FISS) with full sampling grid shown.

Quality control and sampling rate estimations: Following a program of planned FISS expansions from 2014-19, a process of rationialisation of the FISS was undertaken. The goal was to ensure that, given constraints on resources available for implementing the FISS, station selection was such that density indices would be estimated with high precision and low potential for bias. An annual design review process has been developed during which potential FISS designs for the subsequent three years are evaluated according to precision and bias criteria. The resulting proposed designs and their evaluation are presented for review at the June Scientific Review Board (SRB) meetings and potentially modified following SRB input before presentation to the Commissioners at the Work Meeting and Interim Meeting. Annual biological sampling rates for each IPHC Regulatory Area are calculated based on the previous year's catch rates and an annual target of 2000 sampled fish (with 100 additional archive samples).

5.2.2.2 Fishery-independent Trawl Survey (FITS)

The IPHC has participated routinely in the NOAA Fisheries trawl surveys operating in the Bering Sea (Fig. 7, annually since 1998), Aleutian Islands (intermittently since 1997) and Gulf of Alaska (since 1996). The information collected from Pacific halibut caught on these surveys, together with data from the IPHC Fishery-Independent Setline Survey (FISS) and commercial Pacific halibut data, are used directly in estimating indices of abundance and in the stock assessment and to monitor population trends, growth/size, and to supplement understanding of recruitment, distribution, and age composition of young Pacific halibut.




Figure 7. Sampling station design for the 2018 NOAA Bering Sea bottom trawl survey. Black dots are stations sampled in the 2018 "rapid-response" Northern Bering Sea trawl survey and black plus signs are stations sampled in standardized Northern Bering Sea trawl survey.

5.2.2.3 Norton Sound trawl survey

The Alaska Department of Fish and Game's annual Norton Sound trawl survey data contribute to the estimation of Pacific halibut indices of abundance in IPHC Regulatory Area 4E.

5.2.3 Age composition data (both fishery-dependent and fishery-independent)

The annually collected biological samples from commercial fisheries and FISS include otoliths, a crystalline calcium carbonate structure found in the inner ear of fish which growth patterns can be analyzed to estimate the age of fish. Fish age is a key input to stock assessment models that inform management decisions related to fish exploitation. Since inception, the IPHC aged over 1.5 million otoliths manually by trained readers under the stereoscopic microscope.

5.3 Management-supporting information

Successful fisheries management requires rigorous application of the scientific method of problem solving in the development of strategic alternatives and their evaluation on the basis of objectives that integrate ecosystem and human dynamics across space and time into management decision-making (Lane and Stephenson, 1995). This underscores the importance of a holistic understanding of a broad range of factors to deliver on the Commission's objective to develop the stocks of Pacific halibut to the levels that permit the optimum yield from the fishery over time. Management-supporting information beyond IPHC's current research and monitoring programs relate to,



among others, socioeconomic considerations, community development, political constraints, and operational limitations.

Responding to the Commission's "desire for more comprehensive economic information to support the overall management of the Pacific halibut resource in fulfillment of its mandate" (economic study terms of reference adopted at FAC095 and endorsed at AM095 in 2019), between 2019 and 2021 the IPHC conducted a socioeconomic study. The study's core product, Pacific halibut multiregional economic impact assessment (PHMEIA) model, describes economic interdependencies between sectors and regions to bring a better understanding of the role and importance of the Pacific halibut resource to regional economies of Canada and the United States of America (see project report). The model details the within-region production structure of the Pacific halibut sectors (fishing, processing, charter) and cross-regional flows of economic benefits. The model also accounts for economic activity generated through sectors that supply fishing vessels, processing plants, and charter businesses with inputs to production, by embedding Pacific halibut sectors into the model of the entire economy of Canada and the USA. The PHMEIA model fosters stakeholders' better understanding of a broad scope of regional impacts of the Pacific halibut resource. The results highlight that the harvest stage accounts for only a fraction of economic activity that would be forgone if the resource was not available to fishers in the Pacific Northwest. Moreover, the study informs on the vulnerability of communities to changes in the state of the Pacific halibut stock throughout its range, highlighting regions particularly dependent on economic activities that rely on Pacific halibut. Leveraging multiple sources of socioeconomic data, the project provides complementary input for designing policies with desired effects depending on regulators' priorities which may involve balancing multiple conflicting objectives. A good understanding of the localized effects is pivotal to policymakers who are often concerned about community impacts, particularly in terms of impact on employment opportunities and households' welfare.

The economic impact assessment is supplemented by an analysis of the formation of the price paid for Pacific halibut products by final consumers (end-users) that is intended to provide a better picture of Pacific halibut contribution to the gross domestic product (GDP) along the entire value chain, from the hook-to-plate. This supplemental material is available in IPHC's Pacific halibut market analysis.

6. Core focal areas – Planned and opportunistic activities (2022-2026)

Research at IPHC can be classified as "use-inspired basic research" (Stokes 1997) which combines knowledge building with the application of existing and emerging knowledge to provide for the management of Pacific halibut. The four core focal areas: stock assessment, management strategy evaluation, management supporting information, and biology & ecology, all interact with each other as well as with fisheries monitoring activities in the IPHC program of integrated research and monitoring. Progress and knowledge building in one focal area influences and informs application in other core focal areas, also providing insight into future research priorities. The circular feedback loop is similar to the scientific method of observing a problem, creating a hypothesis, testing that hypothesis through research and analysis, drawing conclusions, and refining the hypothesis.

The IPHC Secretariat has been working with IPHC advisory bodies, such as the Scientific Review Board (SRB), and the Commission to conduct scientific research in a way that utilizes the scientific method. Problems are often identified by an advisory body or Commission and hypotheses are developed by the IPHC Secretariat. Research is reviewed by the SRB and refined hypotheses are presented to advisory bodies and the Commission. This process occurs via an annual schedule of meetings, as shown in Fig. 8. In May, an MSE informational session may be held if there is significant progress in the MSE such that it would be useful to prepare stakeholders for the Management Strategy Advisory Board (MSAB) meeting in October. Recommendations related to the MSE, and development of a harvest strategy directed to the Commission are a result of the MSAB meeting. The SRB holds two meetings each year: one in June where requests are typically directed to IPHC Secretariat, and one in September where recommendations are made to the Commission. The June SRB meeting has a focus on research;

the September meeting represents a final check of science products to be presented to the Commission for use in management. The Research Advisory Board (RAB) meets in November to discuss ongoing research, provide guidance and recommend new research projects. The Work Meeting (WM) is held in September and is a working session with IPHC Secretariat and the Commission to prepare for the Interim Meeting (IM) held in November and the Annual Meeting (AM) held in January. Outcomes from the AM include mortality limits (coastwide and by IPHC Regulatory Area), directed fishery season dates, domestic regulations, and requests and recommendations for the IPHC Secretariat. In conjunction with the AM are meetings of the Finance and Administration Committee (FAC), the Conference Board (CB), and the Processor Advisory Board (PAB). The Commission may also hold Special Sessions (SS) throughout the year to take up and make decisions on specific topics.




Figure 8. The typical IPHC annual meeting schedule with the calendar year and fiscal year shown. The meetings, shown in the middle row are: Annual Meeting where the Commission makes many final decisions for that year (AM), an MSE informational session (MSE), Scientific Review Board meetings (SRB), the Commission Work Meeting (WM), the Management Strategy Advisory Board meeting (MSAB), the Research Advisory Board Meeting (RAB), and the Interim Meeting (IM). The annual FISS schedule is also shown.

In addition to the annual meeting process at IPHC, individual core focal areas of research may identify and prioritize research for other core focal areas. For example, stock assessment research often identifies gaps in the knowledge of Pacific halibut biology and ecology, which then identifies priority research for the Biology and Ecology core area. Vice versa, basic biological and ecological research can identify concepts that could be better understood and result in improved implementation in any of the core areas. Furthermore, Management Strategy Evaluation can often be used to identify priority research topics for any core areas by simulation testing to identify research that may have the largest benefit to improving the management of Pacific halibut.

The top priorities of research for various categories in each of the core focal areas are provided below. The top priorities are a subset of the potential research topics in each core focal area. More exhaustive and up-to-date lists of research topics, that may extend beyond a five-year timeframe, can be found in recent meeting documents related to each core focal area.

6.1 Research

6.1.1 Stock Assessment

Within the four assessment research categories, the following topics have been identified as top priorities in order to focus attention on their importance for the stock assessment and management of Pacific halibut. A brief narrative is provided here to highlight the specific use of products from these studies in the stock assessment.

6.1.1.1 Stock Assessment data collection and processing

6.1.1.1.1 Commercial fishery sex-ratio-at-age via genetics

Commercial fishery sex-ratio information has been found to be closely correlated with the absolute scale of the population estimates in the stock assessment and has been identified as the greatest source of uncertainty since 2013. With only four years (2017-20) of commercial sex-ratio-at-age information available for the 2021 stock assessment, the annual genetic assay of fin clips sampled from the landings remains critically important. When the time series grows longer, it may be advantageous to determine the ideal frequency at which these assays need to be conducted. Development of approaches to use archived otoliths, scales or other samples to derive historical estimates (if possible) could provide valuable information on earlier time-periods (with differing fishery and biological properties), and therefore potentially reconcile some of the considerable historical uncertainty in the present stock assessment. This assessment priority directly informs 6.1.3.2 Reproduction as described below.

6.1.1.1.2 Whale depredation accounting and tools for avoidance

Whale depredation currently represents a source of unobserved and unaccounted-for mortality in the assessment and management of Pacific halibut. A logbook program has been phased in over the last several years, in order to record whale interactions observed by commercial harvesters. Estimation of depredation mortality, from logbook records and supplemented with more detailed data and analysis from the FISS represents a first step in accounting for this source of mortality; however, such estimates will likely come with considerable uncertainty. Reduction of depredation mortality through improved fishery avoidance and/or catch protection would be a preferable extension and/or solution to basic estimation. As such, research to provide the fishery with tools to reduce depredation is considered a closely-related high priority. This assessment priority directly informs 6.1.3.4 Mortality and Survival Assessment as described below.

6.1.1.2 Stock Assessment technical development

6.1.1.2.1 Maintaining coordination with the MSE

The stock assessment and MSE operating models have been developed in close coordination, in order to identify plausible hypotheses regarding the processes governing Pacific halibut population dynamics. Important aspects of Pacific halibut dynamics include recruitment (possibly related to extrinsic environmental factors in addition to spawning biomass), size-at-age, movement/migration and spatial patterns in fishery catchability and selectivity. Many approaches developed as part of the tactical stock assessment have been explored in the MSE operating model, and conversely, the MSE operating model has highlighted areas of data uncertainty or alternative hypotheses for exploration in the assessment (e.g. movement rates). Although these two modelling efforts target differing objectives (tactical vs. strategic) continued coordination is essential to ensure that the stock assessment and the MSE represent the Pacific halibut similarly and provide consistent and useful advice for tactical and strategic decision-making.

6.1.1.2.2 Data weighting

The stock assessment currently relies on iterative "Francis" weighting of the age compositional data using a multinomial likelihood formulation (Francis 2011) based on the number of samples available in each year. Exploration of a stronger basis for input sample sizes through analysis of sampling design, estimation of sample weighting and alternative likelihoods may all provide for a more stable approach and a better description of the associated uncertainty.

6.1.1.2.3 Environmental covariates to recruitment

The two long time-series models included in the stock assessment ensemble allow for the Pacific Decadal



Oscillation (PDO; Mantua et al. 1997) to be a binary covariate indicating periods of higher or lower average recruitment. This relationship has been observed to be consistent since its development over 20 years ago (Clark et al 1999) and is re-estimated in each year's stock assessment models. With additional years of data, evaluation of the strength of this relationship, as compared to other metrics of the PDO (e.g., annual deviations, running averages) or other indicators of NE Pacific Ocean productivity should be undertaken in order to provide the best estimates and projections of Pacific halibut recruitment and to provide for alternative hypotheses for use in the MSE. This assessment priority partially informs 6.1.3.2 Reproduction as described below.

6.1.1.2.4 'Leading' parameter estimation

Stock assessments are generally very sensitive to the estimates of leading parameters (stock-recruitment parameters, natural mortality, sex-specific dynamics, etc.). For Pacific halibut some of these are fully integrated into the estimation uncertainty (average unexploited recruitment), or partially integrated (e.g. estimation of natural mortality in two of the four models). As time-series of critically informative data sources like the FISS and the sex-ratio of the commercial landings grow longer it may be possible to integrate additional leading parameters directly in the assessment models and/or include them as nested models within the ensemble.

6.1.1.3 Stock Assessment biological inputs

6.1.1.3.1 Maturity, skip-spawning, and fecundity

Management of Pacific halibut is currently based on reference points that rely on relative female spawning biomass. Therefore, any changes to the understanding of reproductive output – either across age/size (maturity), over time (skip spawning) or as a function of body mass (fecundity) are crucially important. Each of these components directly affects the annual reproductive output estimated in the assessment. Ideally, the IPHC would have a program in place to monitor each of these three reproductive processes over time and use that information in the estimation of the stock-recruitment relationship, and the annual reproductive output relative to reference points. This would reduce the potential for biased time-series estimates created by non-stationarity in these traits (illustrated via sensitivity analyses in several of the recent assessments). However, at present we have only historical time-aggregated estimates of maturity and fecundity schedules. Therefore, the current research priority is to first update our estimates for each of these traits to reflect current environmental and biological conditions. After current stock-wide estimates have been achieved, a program for extending this information to a time-series via transition from research to monitoring can be developed. This assessment priority directly informs 6.1.3.2 Reproduction as described below.

6.1.1.3.2 Stock structure of IPHC Regulatory Area 4B relative to the rest of the convention area

The current stock assessment and management of Pacific halibut assume that IPHC Regulatory Area 4B is functionally connected with the rest of the stock, i.e., that recruitment from other areas can support harvest in Area 4B and that biomass in Area 4B can produce recruits that may contribute to other Areas. Tagging (Webster et al. 2013) and genetic (Drinan et al. 2016) analyses have indicated the potential for Area 4B to be demographically isolated. An alternative to current assessment and management structure would be to treat Area 4B separately from the rest of the coast. This would not likely have a large effect on the coastwide stock assessment as Area 4B represents only approximately 5% of the surveyed stock (Stewart and Webster 2022). However, it would imply that the specific mortality limits for Area 4B could be very important to local dynamics and should be separated from stock-wide trends. Therefore, information on the stock structure for Area 4B has been identified as a top priority. This assessment priority directly informs 6.1.3.1 Migration and Population Dynamics as described below.

6.1.1.3.3 Meta-population dynamics (connectivity) of larvae, juveniles, and adults

The stock assessment and current management procedure treat spawning output, juvenile Pacific halibut abundance, and fish contributing to the fishery yield as equivalent across all parts of the Convention Area. Information on the connectivity of these life-history stages could be used for a variety of improvements to the assessment and current management procedure, including: investigating recruitment covariates, structuring spatial assessment models, identifying minimum or target spawning biomass levels in each Biological Region, refining the stock-recruitment relationship to better reflect source-sink dynamics and many others. Spatial dynamics have been highlighted as a major source of uncertainty in the Pacific halibut assessment for decades and will continue to be of high priority until they are better understood. This assessment priority directly informs 6.1.3.1 Migration and Population Dynamics as described below.

6.1.1.4 Stock Assessment fishery yield

6.1.1.4.1 Biological interactions with fishing gear

In 2020, 16% of the total fishing mortality of Pacific halibut was discarded (Stewart et al. 2021). Discard mortality rates can vary from less than 5% to 100% depending on the fishery, treatment of the catch and other factors (Leaman and Stewart 2017). A better understanding of the biological underpinnings for discard mortality could lead to increased precision in these estimates, avoiding potential bias in the stock assessment. Further, improved biological understanding of discard mortality mechanisms could allow for reductions in this source of fishing mortality, and thereby increased yield available to the fisheries. This assessment priority directly informs 6.1.3.4 Mortality and Survival Assessment as described below.

6.1.1.4.2 Guidelines for reducing discard mortality

Much is already known about methods to reduce discard mortality, in non-directed fisheries as well as the directed commercial and recreational sectors. Promotion and adoption of best handling practices could reduce discard mortality, lead to greater retained yield, and reduce the potential uncertainty associated with large quantities of estimated mortality due to discarding. This assessment priority directly informs 6.1.3.4 Mortality and Survival Assessment as described below.

Outside of the four general assessment categories, the IPHC has recently considered adding close-kin genetics (e.g., Bravington et al. 2016) to its ongoing research program (see section 6.1.3.1). Close-kin mark-recapture can potentially provide estimates of the absolute scale of the spawning output from the Pacific halibut population. This type of information can be fit directly into the stock assessment, and if estimated with a reasonable amount of precision, even a single data point could substantially reduce the uncertainty in the scale of total population estimates. Further, close-kin genetics may provide independent estimates of total mortality (and therefore natural mortality conditioned on catch-at-age), relative fecundity-at-age, and the spatial dynamics of spawning and recruitment. All of these quantities could substantially improve the structure of the current assessment and reduce uncertainty. Data collection of genetic samples from 100% of the sampled commercial landings has been in place since 2017 (as part of the sex-ratio monitoring) and from the FISS since 2021. The genetic analysis required to produce data allowing the estimation of reproductive output and other population parameters from close-kin markrecapture modelling is both complex and expensive, and it could take several years for this project to get fully underway. This five-year plan should consider a pilot evaluation, such that a broader study could be undertaken in the future, providing the likely results would meet the Commission's objectives and prove possible given financial constraints. Research related to close-kin genetics would be pursued under 6.1.3.1 Migration and Population Dynamics as described below.

6.1.2 Management Strategy Evaluation

MSE priorities have been subdivided into three categories: 1) biological parameterisation, 2) fishery parameterization, and 3) technical development. Research provides specifications for the MSE simulations, such as inputs to the Operating Model (OM), but another important outcome of the research is to define the range of plausibility to include in the MSE simulations as a measure of uncertainty. The following topics have been identified as top priorities.

6.1.2.1 MSE Biological and population parameterization

6.1.2.1.1 Distribution of life stages and stock connectivity

Research topics in this category will mainly inform parameterization of movement in the OM, but will also provide further understanding of Pacific halibut movement, connectivity, and the temporal variability. This knowledge may also be used to refine specific MSE objectives to reflect reality and plausible outcomes. Research under Section 6.1.3.1 will inform this MSE priority.

This research includes examining larval and juvenile distribution which is a main source of uncertainty in the OM that is currently not fully incorporated. Outcomes will assist with conditioning the OM, verify patterns simulated from the OM, and provide information to develop reasonable sensitivity scenarios to test the robustness of MPs.

Also included in this number one priority is stock structure research, especially regarding IPHC Regulatory Area 4B. The dynamics of this IPHC Regulatory Area are not fully understood and it is useful to continue research on the connectivity of IPHC Regulatory Area 4B with other IPHC Regulatory Areas.

Finally, genomic analysis of population size is also included in this ranked category because that would help inform development of the OM as well as the biological sustainability objective related to maintaining a minimum spawning biomass in each IPHC Regulatory Area. An understanding of the spatial distribution of population size will help to inform this objective as well as the OM conditioning process.

6.1.2.1.2 Spatial spawning patterns and connectivity between spawning populations

An important parameter that can influence simulation outcomes is the distribution of recruitment across Biological Regions. Continued research in this area will improve the OM and provide justification for parameterising temporal variability. Research includes assigning individuals to spawning areas and establishing temporal and spatial spawning patterns. Outcomes may also provide information on recruitment strength and the relationship with environmental factors. For example, recent work by Sadorus et al (2020) used a biophysical and spatio-temporal models to examine connectivity across the Bering Sea and Gulf of Alaska. Furthermore, close-kin mark-recapture (Bravington et al. 2016) may provide insights into spatial relationships between juveniles and adults as well as abundance in specific regions. Research under Sections 6.1.3.1 and 6.1.3.2 will inform this MSE priority.

6.1.2.1.3 Understanding growth variation

Changes in the average weight-at-age of Pacific halibut is one of the major drivers of changes in biomass over time. The OM currently simulates temporal changes in weight-at-age via a random autocorrelated process which is unrelated to population size or environmental factors. Ongoing research in drivers related to growth in Pacific halibut will help to improve the simulation of weight-at-age. Research under Section 6.1.3.3 will inform this MSE priority.

6.1.2.1.4 MSE fishery parameterization

The specifications of fisheries and their parameterizations involved consultation with Pacific halibut stakeholders but some aspects of those parameterizations benefit from targeted research. One specific example is knowledge of discarding and discard mortality rates in directed and non-directed fisheries. Discard mortality can be a significant source of fishing mortality in some IPHC Regulatory Areas and appropriately modelling that mortality will provide a more robust evaluation of MPs. Research under Sections 6.1.3.4 and 6.1.3.5 will inform this MSE priority.

6.1.2.2 MSE technical development

Technical improvements to the MSE framework will allow for rapid development of alternative operating models and efficient simulation of management strategies for future evaluation. Coordination with the technical development of the stock assessment (Section 6.1.1.2.1) is necessary to ensure consistent assumptions and hypotheses for tactical (i.e. stock assessment) and strategic (i.e. MSE) models. Investigations done in the stock assessment will inform the stock assessment, which will then be informed by investigations using the closed-loop simulation framework. Multi-year assessments may allow for additional opportunity to coordinate between stock assessment and MSE.

6.1.2.2.1 Alternative migration scenarios

Including alternative migration hypotheses in the MSE simulations will assist in identifying management procedures that are robust to this uncertainty. This exploration will draw on general research on the movement and migration of Pacific halibut, observations from FISS and fisheries data, and outcomes of the stock assessment. Identification of reasonable hypotheses for the movement of Pacific halibut is essential to the robust investigation of management procedures. Research under Section 6.1.3.1 will inform this MSE priority.

6.1.2.2.2 Realistic simulations of estimation error

Closed loop simulation uses feedback from the management procedure to update the population in the projections. The management procedure consists of data collection, an estimation model, and harvest rules; currently IPHC uses a stock assessment as the estimation model. Future development of an efficient simulation process to mimic the stock assessment will more realistically represent the current management process. This involves using multiple estimation models to represent the ensemble and appropriately adding data and updating those models in the simulated projections. Improvements to the current MSE framework include adding additional estimation models to better represent the ensemble stock assessment, ensuring that the simulated estimation accurately represent the stock assessment now and, in the future, and speeding up the simulation process.

6.1.2.2.3 Incorporate additional sources of implementation uncertainty

Implementation uncertainty consists of three subcategories: 1) decision-making uncertainty, 2) realized uncertainty, and 3) perceived uncertainty. Decision-making uncertainty is the difference between mortality limits determined from the management procedure and those adopted by the Commission. This uncertainty is currently not implemented in the MSE framework but has been requested by the SRB and the independent peer review of the MSE. Realized uncertainty is the difference between the mortality limit set by the Commission and the actual mortality realized by the various fisheries. This type of uncertainty is currently partially implemented in the MSE framework. Finally, perceived uncertainty is the difference between the realized mortality and the estimated mortality limits from the various fisheries, which would be used in the estimation model. This third type of implementation uncertainty has not been implemented in the MSE framework. Implementing decision-making uncertainty is a priority for the MSE and will assist in understanding the performance of management procedures when they may not be followed exactly.

6.1.2.3 MSE Program of Work for 2021–2023

Following the 11th Special Session of the IPHC, an MSE program of work for 2021–2023 was developed. Seven tasks were identified that pertained to further developments of the MSE framework, evaluation of alternative MPs, and improvements in evaluation and presentation of results. <u>Table 1</u> lists these tasks and provides a brief description. Additional details can be found in the program of work available on the MSE webpage.

Table 1. Tasks recommended by the Commission at SS011 (<u>IPHC-2021-SS011-R</u> para 7) for inclusion in the IPHC Secretariat MSE Program of Work for 2021–23.

ID	Category	Task	Deliverable
F.1	Framework	Develop migration scenarios	Develop OMs with alternative migration scenarios
F.2	Framework	Implementation variability	Incorporate additional sources of implementation variability in the framework
F.3	Framework	Develop more realistic simulations of estimation error	Improve the estimation model to more adequately mimic the ensemble stock assessment
F.5	Framework	Develop alternative OMs	Code alternative OMs in addition to the one already under evaluation.
M.1	MPs	Size limits	Identification, evaluation of size limits
M.3	MPs	Multi-year assessments	Evaluation of multi-year assessments
E.3	Evaluation	Presentation of results	Develop methods and outputs that are useful for presenting outcomes to stakeholders and Commissioners

6.1.2.4 Potential Future MSE projects

Management Strategy Evaluation is an iterative process where new management procedures may be evaluated, current management procedures may be re-evaluated under different assumptions, and the understanding of the population, environment, and fisheries may be updated with new information stemming from the stock assessment and biological/ecological research. The current Program of Work (<u>Table 1</u>) focuses on two elements of Management Procedures, but in the future other elements may be of interest, such as distribution procedures. The research being done now will inform the development of the MSE in the future to ensure a robust evaluation of any management procedure.

6.1.3 Biology and Ecology

Capitalizing on the outcomes of the previous 5-year plan (IPHC–2019–BESRP-5YP) (<u>Appendix I</u>), the IPHC Secretariat has identified five research areas that will provide key inputs for stock assessment and the MSE process. In addition to linking genetics and genomics with migration and distribution studies in the newly coined area of Migration and Population Dynamics, the IPHC Secretariat has incorporated a novel research area on Fishing Technology. A series of key objectives for each of the five research areas have been identified that integrate with specific needs for stock assessment and MSE processes and that are ranked according to their relevance (<u>Appendix II</u>). To further describe the IPHC Secretariat's rationale for establishing research priorities, a ranked list of biological uncertainties and parameters for stock assessment and the MSE process and their links to research activities and outcomes derived from the IPHC 5-Year Program of Integrated Research and Monitoring (2022-2026) are provided in <u>Appendix III</u> and <u>Appendix IV</u>.

6.1.3.1 Migration and Population Dynamics

Genetic and genomic studies aimed at improving current knowledge of Pacific halibut migration and population dynamics throughout all life stages in order to achieve a complete understanding of stock structure and distribution across the entire distribution range of Pacific halibut in the North Pacific Ocean and the biotic and abiotic factors that influence it (specifically excluding satellite tagging). Specific objectives in this area include:

- Improve current knowledge of the genetic structure of the Pacific halibut population through the use of state-of-the-art low-coverage whole genome resequencing approaches. Establishment of genetic signatures of spawning sites.
- Improve our understanding of the mechanisms and magnitude of larval connectivity in the North Pacific Ocean. Identification of environmental and biological predictors of larval abundance and recruitment.
- Improve our understanding of spawning site contributions to nursery/settlement areas in relation to yearclass, recruit survival and strength, and environmental conditions in the North Pacific Ocean. Measure of genetic diversity of Pacific halibut juveniles from the eastern Bering Sea and the Gulf of Alaska.
- Improve our understanding of the relationship between nursery/settlement origin and adult distribution and abundance over temporal and spatial scales. Genomic assignment of individuals to source populations and assessment of distribution changes.
- Integrate analyses of Pacific halibut connectivity and distribution changes by incorporating genomic approaches.
- Improve estimates of population size, migration rates among geographical regions, and demographic parameters (e.g. fecundity-at-age, survival rate), through the application of close-kin mark-recapture-based approaches.
- Improve our understanding of the influences of oceanographic and environmental variation on connectivity, population structure and adaptation at a genomic level using seascape genomics approaches.
- Exploration and development of alternative methods for aging Pacific halibut based on genetic analyses of DNA methylation patterns in tissues (fin clips).
- Exploration of methods for individual identification based on computer-assisted tail image matching systems as an alternative for traditional mark and recapture tagging.

6.1.3.2 Reproduction

Studies aimed primarily at addressing two critical issues for stock assessment analysis based on estimates of female spawning biomass: 1) the sex ratio of the commercial catch and 2) maturity estimations. Specific objectives in this area include:

- Continued improvement of genetic methods for accurate sex identification of commercial landings from fin clips and otoliths in order to incorporate recent and historical sex-at-age information into the stock assessment process.
- Improve our understanding of the temporal progression of reproductive development and gamete production during an entire annual reproductive cycle in female and male Pacific halibut.
- Update current maturity-at-age estimates.
- Provide estimates of fecundity-at-age and fecundity-at-size.
- Investigate the possible presence of skip spawning in Pacific halibut females.

- Improve accuracy in current staging criteria of maturity status used in the field.
- Investigate possible environmental effects on the ontogenetic establishment of the phenotypic sex and their influence on sex ratios in the adult Pacific halibut population.
- Improve our understanding of potential temporal and spatial changes in maturity schedules and spawning patterns in female Pacific halibut and possible environmental influences.
- Improve our understanding of the genetic basis of variation in age and/or size-at-maturity, fecundity, and spawning timing, by conducting genome-wide association studies.

6.1.3.3 Growth

Studies aimed at describing the role of factors responsible for the observed changes in size-at-age and at evaluating growth and physiological condition in Pacific halibut. Specific objectives in this area include:

- Evaluate possible variation in somatic growth patterns in Pacific halibut as informed by physiological growth markers, physiological condition, energy content and dietary influences.
- Investigate the effects of environmental and ecological conditions that may influence somatic growth in Pacific halibut. Evaluate the relationship between somatic growth and temperature and trophic histories in Pacific halibut through the integrated use of physiological growth markers.
- Improve our understanding of the genetic basis of variation in somatic growth and size-at-age by conducting genome-wide association studies.

6.1.3.4 Mortality and Survival Assessment

Studies aimed at providing updated estimates of discard mortality rates (DMRs) for Pacific halibut in the guided recreational fisheries and at evaluating methods for reducing mortality of Pacific halibut. Specific objectives in this area include:

- Provide information on the types of fishing gear and fish handling practices used in the Pacific halibut recreational (charter) fishery as well as on the number and size composition of discarded Pacific halibut in this fishery.
- Establish best handling practices for reducing discard mortality of Pacific halibut in recreational fisheries.
- Investigate new methods for improved estimation of depredation mortality from marine mammals.

6.1.3.5 Fishing Technology

Studies aimed at developing methods that involve modifications of fishing gear with the purpose of reducing Pacific halibut depredation and bycatch. Specific objectives in this area include:

- Investigate new methods for whale avoidance and/or deterrence for the reduction of Pacific halibut depredation by whales (e.g. catch protection methods).
- Investigate physiological and behavioral responses of Pacific halibut to fishing gear in order to reduce bycatch.

6.2 Monitoring

The Commission's extensive monitoring programs include both direct data collection and coordination with domestic agencies to produce both fishery-dependent and fishery-independent information on the stock and fishery trends, and other information. These critical sources include estimates of fishing mortality from all

fisheries encountering Pacific halibut, biological sampling from these fisheries as well as catch-rates and biological sampling from longline and trawl surveys. Monitoring data provide the basis for stock assessment and MSE analysis, many biological research studies, and some inputs directly to the decision-making process (Figure 4). While not the primary focus of this 5-year plan, a basic summary of the components led by the IPHC and those that are provided by domestic agencies is provided below.

6.2.1 Fishery-dependent data

Data collection and monitoring activities aimed at providing standardised time-series of mortality, fishery, and biological data from both direct target fisheries as well as fisheries that incidentally catch Pacific halibut. Directed commercial fisheries data are managed by IPHC. Non-directed commercial discard mortality data, subsistence fisheries data, and recreational fisheries data are managed by Contracting Party domestic agencies.

6.2.1.1 Directed commercial fisheries data

6.2.1.2 Annually review the spatial distribution of sampling effort among ports, data collection methods, sampling rates, and quality assurance/quality control (QAQC) processes, including in-season review of port sampling activities

Ensure current data collection efforts meet current and future needs of stock assessment, MSE and management. Collaborate and coordinate with other Secretariat functions to develop methods and procedures for incorporating promising research results into long-term monitoring program. The IPHC relies on domestic and Tribal agency programs to report annual mortality from incidental catches in non-directed commercial fisheries, catches from subsistence fisheries, and catches from recreational fisheries. Non-directed commercial discard mortality data

Annually collaborate with observer programs and other partners to ensure robust data collection and sampling, QAQC processes, and reporting of incidental catch and mortality, as well as biological sampling.

6.2.1.3 Subsistence fisheries data

Annually collaborate with Tribal, State and Federal agencies of each Contracting Party to ensure high quality data collection, sampling, and reporting in the subsistence fisheries in Canada and the United States of America.

6.2.1.4 Recreational fisheries data

Annually collaborate with National/State agencies of each Contracting Party to ensure and validate high quality data and reporting of recreational fishery mortality estimates and biological data.

6.2.2 Fishery-independent data

Data collection and monitoring activities aimed at providing a standardised time-series of biological and ecological data that is independent of the fishing fleet.

6.2.2.1 Fishery-independent setline survey (FISS)

An annual review process for the FISS station design has been developed (Fig. 9) and is expected to continue in coming years. This process involves scientific review of proposed FISS designs by the Scientific Review Board and includes input from stakeholders prior to review and approval of designs by the Commissioners.

Direct weighing of Pacific halibut has been integrated into the annual FISS sampling since 2019 and will continue into the future to ensure accurate estimation of WPUE and other weight-derived quantities. Sample rates for genetic monitoring will need to be determined for future sampling. Sampling rates of otoliths for aging, archive otoliths and tagged fish will continue to be reviewed annually to ensure the data needs of the IPHC stock assessment and research program are met. Annual FISS sampler training and data QAQC (including at point of

data collection and during post-sampling review) will ensure high quality data from the FISS program. Procedures are reviewed annually.




Figure 9. Timeline of annual FISS design review process.

6.2.2.2 Fishery-independent Trawl Survey (FITS)

The IPHC will continue to collaborate with NMFS on sampling procedures for Pacific halibut on the placement of an IPHC sampler onboard a survey vessel for the collection of biological data.

6.2.3 Age composition data (both fishery-dependent and fishery-independent)

The IPHC Secretariat is looking at options for supplementing current Pacific halibut ageing protocol with automatized ageing that does not require extensive otolith-reader training. The IPHC is investigating the potential use of artificial intelligence (AI) for determining the age of Pacific halibut from images of collected otoliths. The Secretariat is in the process of initializing creation of a database of pictures with expert-provided labels, utilizing previously aged otoliths, and assessing the option for the development of a Convolutional Neural Network (CNN) model specifically designed for image classification to determine Pacific halibut age. The goal is to create an AI-based age determination system that complements traditional methods for reliable fish stock assessment and management advice.

6.3 Potential of integrating human dynamics into management decision-making

The evolution of modern fisheries management is taking a transformative turn, emphasizing the integration of human dynamics into decision-making processes. As our world becomes more interconnected through globalization, understanding the intricate human dimension of the fisheries sector is emerging as a critical aspect of sustainable resource management. This forward-looking approach seeks to proactively address challenges while capitalizing on new opportunities.

In a global marketplace where local and imported products compete for consumer attention, vulnerability to disruptions, as evidenced by the COVID-19 pandemic (OECD 2020), has highlighted the need for adaptable strategies embracing the broader picture encompassing external influences. Recent IPHC's socioeconomic study underlines the far-reaching impacts of such dynamics, showcasing the income fluctuations experienced by

households dependent on Pacific halibut during the pandemic. Acknowledging these complexities, there is a growing realization of the need for expanding the scope of management-supporting information the IPHC provides beyond stock condition.

The question of how small remote communities can capitalize on the high prices that the final customers are paying for premium seafood products demands innovative thinking. In 2021, fresh Alaskan Pacific halibut fillets routinely sold for USD 24-28 a pound, and often more, in downtown Seattle (e.g. USD 38 at Pike Place Market). Pacific halibut dishes at the restaurants typically sell for USD 37-43 for a dish including a 6oz fish portion. The IPHC's socioeconomic study detailed the geography of impacts of the Pacific halibut fisheries, providing a coherent picture of the exposure of fisheries-dependent households by location to changes in resource availability, but paying closer attention to quantifying leakage of economic benefits from communities strongly involved in fisheries, highlighted that the local earnings often do not align with how much fishing occurs within the community. This suggests the need for research focused on how to operationalize social equity in the context of the globalized market dynamics and the pursuit of stock sustainability.

In parallel, the accelerating impacts of climate change is placing fisheries at the forefront of environmental challenges. The rapid increase in water temperature off the coast of Alaska in 2014-16, termed *the blob*, exemplifies the changes that disrupt ecosystems and fisheries (Cheung and Frölicher 2020), and may have a long-term impact on Pacific halibut distribution. The consequences may include shifts in the distribution of benefits, but possibly go further, affecting the stability of agreements over allocation of a shared resource. Research on decision quality under fast-progressing climate-induced changes to stock distribution emerges as an avenue for impactful work.

Conflicting objectives among stakeholders regarding the use of limited resource in the context of globalization, calls for social equity and climate change are a major challenge of decision-making in fisheries management. Integrating approaches aimed at understanding the human dynamics and external factors with stock assessment and MSE can assist fisheries in bridging the gap between the current and the optimal performance without compromising the stock biological sustainability. For example, socioeconomic performance metrics presented alongside already developed biological/ecological performance metrics would supplement IPHC's portfolio of tools for assessing policy-oriented issues (as requested by the Commission, IPHC-2021-AM097-R, AM097-Req.02) and support decision-making. Moreover, continuing investment in understanding the human dimension of Pacific halibut fishing can also inform on other drivers such as human behavior or human organization that affect the dynamics of fisheries, and thus contribute to improved accuracy of the stock assessment and the MSE (Lynch et al.2018). As such, it can contribute to research integration at the IPHC and provide a complementary resource for the development of harvest control rules.

Lastly, Pacific halibut value is also in its contribution to the diet through subsistence fisheries and importance to the traditional users of the resource. To native people, traditional fisheries constitute a vital aspect of local identity and a major factor in cohesion. One can also consider the Pacific halibut's existence value as an iconic fish of the Pacific Northwest. Recognizing and adopting such an all-encompassing definition of the Pacific halibut resource contribution, the IPHC echoes a broader call to include the human dimension into the research on the impact of management decisions, as well as changes in environmental or stock conditions.

7. Amendment

The intention is to ensure the plan is kept as a 'living plan', that is reviewed and updated annually based on the resources available to undertake the work of the Commission (e.g. internal and external fiscal resources, collaborations, internal expertise). The IPHC Secretariat is committed to ensuring an exceptional level of transparency and commitment to the principles of open science.

8. References

- Bravington, M.V., Skaug, H.J., and Anderson, E.C. 2016. Close-Kin Mark-Recapture. Statistical Science 31(2): 259-274. doi:10.1214/16-sts552.
- Cheung, William W.L., and Thomas L. Frölicher. 2020. "Marine Heatwaves Exacerbate Climate Change Impacts for Fisheries in the Northeast Pacific." Scientific Reports 10 (1): 1–10. https://doi.org/10.1038/s41598-020-63650-z.
- Clark, W.G., Hare, S.R., Parma, A.M., Sullivan, P.J., and Trumble, R.J. 1999. Decadal changes in growth and recruitment of Pacific halibut (*Hippoglossus stenolepis*). Canadian Journal of Fisheries and Aquatic Sciences 56: 242-252.
- Drinan, D.P., Galindo, H.M., Loher, T., and Hauser, L. 2016. Subtle genetic population structure in Pacific halibut Hippoglossus stenolepis. J Fish Biol 89(6): 2571-2594. doi:10.1111/jfb.13148.
- Francis, R.I.C.C. 2011. Data weighting in statistical fisheries stock assessment models. Canadian Journal of Fisheries and Aquatic Sciences 68: 1124-1138.
- Fry, Cristy. 2021. "Seawatch: Halibut Prices Soaring." Homer News, 2021. https://www.homernews.com/business/seawatch-halibut-prices-soaring/.
- Lane, Daniel E, and R L Stephenson. 1995. "Fisheries Management Science: The Framework to Link Biological, Economic, and Social Objectives in Fisheries Management." Aquatic Living Resources 8 (3): 215–21.
- Leaman, B.M., and Stewart, I.J. 2017. 2.12 Research basis for estimated discard mortality rates used for Pacific halibut in longline and trawl fisheries. IPHC Report of Assessment and Research Activities 2016: 133-172.
- Lynch, Patrick D, Richard D Methot, and Jason S Link. 2018. "Implementing a Next Generation Stock Assessment Enterprise: Policymakers' Summary." NOAA Technical Memorandum NMFS-F/SPO-183.
- OECD. 2020. "Fisheries, Aquaculture and COVID-19: Issues and Policy Responses." Tackling Coronavirus (Covid-19).
- Mantua, N.J., Hare, S.R., Zhang, Y., Wallace, J.R., and Francis, R.C. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. Bulletin of the American Meteorological Society 78(6): 1069-1079.
- Sadorus, Lauri L., Esther D. Goldstein, Raymond A. Webster, William T. Stockhausen, Josep V. Planas, and Janet T. Duffy-Anderson. 2020. "Multiple life-stage connectivity of Pacific halibut (*Hippoglossus stenolepis*) across the Bering Sea and Gulf of Alaska." Fisheries Oceanography 30 (2): 174-193. https://doi.org/10.1111/fog.12512
- Stewart, I., and Webster, R. 2022. Overview of data sources for the Pacific halibut stock assessment, harvest policy, and related analyses. IPHC-2022-SA-02. 56 p.
- Stewart, I., Hicks, A., Webster, R., and Wilson, D. 2021. Stock assessment: Summary of the data, stock assessment, and harvest decision table for Pacific halibut (Hippoglossus stenolepis) at the end of 2020. IPHC-2021-AM097-08. 19 p.
- Webster, R.A., Clark, W.G., Leaman, B.M., and Forsberg, J.E. 2013. Pacific halibut on the move: a renewed understanding of adult migration from a coastwide tagging study. Canadian Journal of Fisheries and Aquatic Sciences 70(4): 642-653. doi:10.1139/cjfas-2012-0371.



ACKNOWLEDGEMENTS

This updated document was developed by Dr. David Wilson, Dr. Josep Planas, Dr. Ian Stewart, Dr. Allan Hicks, Dr. Ray Webster, and Dr. Basia Hutniczak, in collaboration with other current members of the IPHC Secretariat.

APPENDICES

Appendix I: Outcomes of the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21)

Appendix II: Biological research areas in the 5-Year Program of Integrated Research and Monitoring (2022-

2026) and ranked relevance for stock assessment and management strategy evaluation

Appendix III: List of ranked research priorities for stock assessment

Appendix IV: List of ranked research priorities for management strategy evaluation

Appendix V: Proposed schedule of outputs

Appendix VI: Proposed schedule with funding and staffing indicators

APPENDIX I

Outcomes of the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21) (IPHC-2019-BESRP-5YP)

A. Outcomes by Research Area:

1. Migration and Distribution.

1.1. <u>Larval and juvenile connectivity and early life history studies</u>. Planned research outcomes: improved understanding of larval and juvenile distribution.

Main results:

- Larval connectivity between the Gulf of Alaska and the Bering Sea occurs through large island passes across the Aleutian Island chain.
- The degree of larval connectivity between the Gulf of Alaska and the Bering Sea is influenced by spawning location.
- Spawning locations in the western Gulf of Alaska significantly contribute Pacific halibut larvae to the Bering Sea.
- Pacific halibut juveniles counter-migrate from inshore settlement areas in the eastern Bering Sea into the Gulf of Alaska through Unimak Pass.
- Elemental signatures of otoliths from juvenile Pacific halibut vary geographically at a scale equivalent to IPHC regulatory areas.

Publications:

Sadorus, L.; Goldstein, E.; Webster, R.; Stockhausen, W.; Planas, J.V.; Duffy-Anderson, J. Multiple life-stage connectivity of Pacific halibut (*Hippoglossus stenolepis*) across the Bering Sea and Gulf of Alaska. *Fisheries Oceanography*. 2021. 30:174-193. doi: https://doi.org/10.1111/fog.12512.

Loher, T., Bath, G. E., Wischniowsky, S. The potential utility of otolith microchemistry as an indicator of nursery origins in Pacific halibut (*Hippoglossus stenolepis*) in the eastern Pacific: the importance of scale and geographic trending. *Fisheries Research*. 2021. 243: 106072. https://doi.org/10.1016/j.fishres.2021.106072.

Links to 5-Year Research Plan (2022-2026):

- Evaluate the level of genetic diversity among juvenile Pacific halibut in the Gulf of Alaska and the Bering sea due to admixture.
- Assignment of individual juvenile Pacific halibut to source populations.

<u>Integration with Stock Assessment and MSE</u>: The relevance of research outcomes from activities in this research area for stock assessment is in the improvement of estimates of productivity. Research outcomes will be used to generate potential recruitment covariates and to inform minimum spawning biomass targets by Biological Region and represent one of the top three biological inputs into stock assessment. The relevance of these research outcomes for MSE is in the improvement of the parametrization of the Operating Model and represent the top ranked biological input into the MSE.



2. Reproduction.

2.1 Sex ratio of commercial landings. Planned research outcomes: sex ratio information.

Main results:

- Establishment of TaqMan-based genetic assays for genotyping Pacific halibut in the IPHC Biological Laboratory.
- Sex ratio information for the 2017-2020 commercial landings.
- Transfer of genotyping efforts for sex identification to IPHC monitoring program.

Links to 5-Year Research Plan (2022-2026):

- Monitoring effort.
- Histological maturity assessment. Planned research outcomes: updated maturity schedule.

Main results:

- Oocyte developmental stages have been characterized and fully described in female Pacific halibut for the first time.
- Oocyte developmental stages have been used for the classification of female developmental stages and to be able to characterize female Pacific halibut as group synchronous with determinate fecundity.
- Female developmental stages have been used for the classification of female reproductive phases and to be able to characterize female Pacific halibut as following an annual reproductive cycle with spawning in January and February.
- Female developmental stages and reproductive phases of females collected in the central Gulf of Alaska have been used to identify the month of August as the time of the transition between the Vtg2 and Vtg3 developmental stages marking the beginning of the spawning capable reproductive phase.
- Future gonad collections for revising maturity schedules and estimating fecundity can be conducted in August during the FISS.

Publications:

- Fish, T., Wolf, N., Harris, B.P., Planas, J.V. A comprehensive description of oocyte developmental stages in Pacific halibut, *Hippoglossus stenolepis*. *Journal of Fish Biology* 2020. 97: 1880-1885. doi: 10.1111/jfb.14551.
- Fish, T., Wolf, N., Smeltz, T. S., Harris, B. P., and Planas, J. V. Reproductive Biology of Female Pacific Halibut (Hippoglossus stenolepis) in the Gulf of Alaska. Frontiers in Marine Science 2022. 9:801759. doi: 10.3389/fmars.2022.801759.

Links to 5-Year Research Plan (2022-2026):

• Revision of maturity schedule by gonad collection during the FISS, as informed by previous studies on reproductive development.

• Estimation of fecundity by age and size, as informed by previous studies demonstrating determinate fecundity.

Integration with Stock Assessment and MSE: Research activities in this Research Area aim at providing information on key biological processes related to reproduction in Pacific halibut (maturity and fecundity) and to provide sex ratio information of Pacific halibut commercial landings. The relevance of research outcomes from these activities for stock assessment is in the scaling of Pacific halibut biomass and in the estimation of reference points and fishing intensity. These research outputs will result in a revision of current maturity schedules and will be included as inputs into the stock assessment and represent the most important biological inputs for stock assessment. The relevance of these research outcomes for MSE is in the improvement of the simulation of spawning biomass in the Operating Model.

3. Growth.

3.1 <u>Identification of physiological growth markers and their application for growth pattern evaluation</u>. Planned research outcomes: informative physiological growth markers.

Main results:

- Transcriptomic profiling by RNAseq of white skeletal muscle from juvenile Pacific halibut subjected to growth suppression and to growth stimulation resulted in the identification of a number of genes that change their expression levels in response to growth manipulations.
- Proteomic profiling by LC-MS/MS of white skeletal muscle from juvenile Pacific halibut subjected to growth suppression and to growth stimulation resulted in the identification of a number of proteins that change their abundance in response to growth manipulations.
- Genes and proteins that changed their expression levels in accordance to changes in the growth rate in juvenile Pacific halibut were selected as putative growth markers for future studies on growth pattern evaluation.

Publications:

Planas et al. 2022. In Preparation.

Links to 5-Year Research Plan (2022-2026):

- Application of identified growth markers in studies aiming at investigating environmental
 influences on growth patterns and at investigating dietary influences on growth patterns and
 physiological condition.
- 3.2 <u>Environmental influences on growth patterns</u>. Planned research outcomes: information on growth responses to temperature variation.

Main results:

• Laboratory experiments under controlled temperature conditions have shown that temperature affects the growth rate of juvenile Pacific halibut through changes in the expression of genes that regulate growth processes.

Publications:

Planas et al. 2022. In Preparation.

Links to 5-Year Research Plan (2022-2026):

- Identification of temperature-specific responses in skeletal muscle through comparison between transcriptomic responses to temperature-induced growth changes and to density- and stress-induced growth changes.
- Application of growth markers for additional studies investigating the link between environmental variability and growth patterns and the effects of diet (prey quality and abundance) on growth and physiological condition.

<u>Integration with Stock Assessment and MSE</u>: Research activities conducted in this Research Area aim at providing information on somatic growth processes driving size-at-age in Pacific halibut. The relevance of research outcomes from these activities for stock assessment resides, first, in their ability to inform yield-per-recruit and other spatial evaluations for productivity that support mortality limit-setting, and second, in that they may provide covariates for projecting short-term size-at-age and may help delineate between fishery and environmental effects, thereby informing appropriate management responses. The relevance of these research outcomes for MSE is in the improvement of the simulation of variability and to allow for scenarios investigating climate change.

4. Mortality and Survival Assessment.

4.1 <u>Discard mortality rate estimation in the longline Pacific halibut fishery</u>. Planned research outcomes: experimentally-derived DMR.

Main results:

- Different hook release methods used in the longline fishery result in specific injury profiles and viability classification.
- Plasma lactate levels are high in Pacific halibut with the lowest viability classification.
- Mortality of discarded fish with the highest viability classification is estimated to be between 4.2 and 8.4%.

Publications:

- Kroska, A.C., Wolf, N., Planas, J.V., Baker, M.R., Smeltz, T.S., Harris, B.P. Controlled experiments to explore the use of a multi-tissue approach to characterizing stress in wild-caught Pacific halibut (*Hippoglossus stenolepis*). *Conservation Physiology* 2021. 9(1):coab001; doi:10.1093/conphys/coab001.
- Loher, T., Dykstra, C.L., Hicks, A., Stewart, I.J., Wolf, N., Harris, B.P., Planas, J.V. Estimation of postrelease longline mortality in Pacific halibut using acceleration-logging tags. *North American Journal of Fisheries Management*. 2022. 42: 37-49. DOI: https://doi.org/10.1002/nafm.10711.

Links to 5-Year Research Plan (2022-2026):

- Integration of information on capture and handling conditions, injury and viability assessment and physiological condition will lead to establishing a set of best handling practices in the longline fishery.
- 4.2 Discard mortality rate estimation in the guided recreational Pacific halibut fishery. Planned research outcomes: experimentally-derived DMR.

Main results:

- Field experiments testing two different types of gear types (i.e. 12/0 and 16/0 circle hooks) resulted in the capture, sampling and tagging of 243 Pacific halibut in IPHC Regulatory Area 2C (Sitka, AK) and 118 in IPHC Regulatory Area 3A (Seward, AK).
- The distributions of fish lengths by regulatory area and by hook size were similar.

Links to 5-Year Research Plan (2022-2026):

- Estimation of discard mortality rate in the guided recreational fishery.
- Integration of information on capture and handling conditions, injury and viability assessment and physiological condition linked to survival.
- Establishment of a set of best handling practices in the guided recreational fishery.

<u>Integration with Stock Assessment and MSE</u>: The relevance of research outcomes from these activities for stock assessment resides in their ability to improve trends in unobserved mortality in order to improve estimates of stock productivity and represent the most important inputs in fishery yield for stock assessment. The relevance of these research outcomes for MSE is in fishery parametrization

5. Genetics and genomics.

<u>5.1</u> <u>Generation of genomic resources for Pacific halibut</u>. Planned research outcomes: sequenced genome and reference transcriptome.

Main results:

- A first draft of the chromosome-level assembly of the Pacific halibut genome has been generated.
- The Pacific halibut genome has a size of 602 Mb and contains 24 chromosome-size scaffolds covering 99.8% of the complete assembly with a N50 scaffold length of 27 Mb at a coverage of 91x.
- The Pacific halibut genome has been annotated by NCBI and is available as NCBI Hippoglossus stenolepis Annotation Release 101 (https://www.ncbi.nlm.nih.gov/assembly/GCA 022539355.2/).
- Transcriptome (i.e. RNA) sequencing has been conducted in twelve tissues in Pacific halibut and
 the raw sequence data have been deposited in NCBI's Sequence Read Archive (SRA) under the
 bioproject number PRJNA634339 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA634339)
 and with SRA accession numbers SAMN14989915 SAMN14989926.

Publications:

Jasonowicz, A.C., Simeon, A., Zahm, M., Cabau, C., Klopp, C., Roques, C., Iampietro, C., Lluch, J., Donnadieu, C., Parrinello, H., Drinan, D.P., Hauser, L., Guiguen, Y., Planas, J.V. Generation of a chromosome-level genome assembly for Pacific halibut (*Hippoglossus stenolepis*) and characterization of its sex-determining genomic region. *Molecular Ecology Resources*. 2022. In *Press*. doi: https://doi.org/10.1111/1755-0998.13641.

Jasonowicz et al. 2022. In Preparation.

Links to 5-Year Research Plan (2022-2026):

• Genome-wide analysis of stock structure and composition.

5.2 <u>Determine the genetic structure of the Pacific halibut population in the Convention Area</u>. Planned research outcomes: genetic population structure.

Main results:

- The collection of winter genetic samples in the Aleutian Islands completed the winter sample collection needed to conduct studies on the genetic population structure of Pacific halibut in the Convention Area.
- Initial results of low coverage whole genome resequencing of winter samples indicate that an average of 26.5 million raw sequencing reads per obtained per sample that provided average individual genomic coverages for quality filtered alignments of 3.2x.

Links to 5-Year Research Plan (2022-2026):

• Fine-scale delineation of population structure, with particular emphasis on IPHC Regulatory 4B structure.

Integration with Stock Assessment and MSE: The relevance of research outcomes from these activities for stock assessment resides in the introduction of possible changes in the structure of future stock assessments, as separate assessments may be constructed if functionally isolated components of the population are found (e.g. IPHC Regulatory Area 4B), and in the improvement of productivity estimates, as this information may be used to define management targets for minimum spawning biomass by Biological Region. These research outcomes provide the second and third top ranked biological inputs into stock assessment. Furthermore, the relevance of these research outcomes for MSE is in biological parametization and validation of movement estimates and of recruitment distribution.



B. List of ranked biological uncertainties and parameters for stock assessment (SA) and their links to research areas and activities contemplated in the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21)

SA Rank	Research outcomes	Relevance for stock assessment	Specific analysis input	Research Area	Research activities
	Updated maturity schedule		Will be included in the stock assessment, replacing the current schedule last updated in 2006		Histological maturity assessment
Biological	Incidence of skip spawning	Scale biomass and	Will be used to adjust the asymptote of the maturity schedule, if/when a time-series is available this will be used as a direct input to the stock assessment		Examination of potential skip spawning
input	Fecundity-at-age and -size information	reference point estimates	Will be used to move from spawning biomass to egg-output as the metric of reproductive capability in the stock assessment and management reference points	Reproduction	Fecundity assessment
	Revised field maturity classification		Revised time-series of historical (and future) maturity for input to the stock assessment		Examination of accuracy of current field macroscopic maturity classification
2. Biological	Stock structure of IPHC Regulatory Area 4B relative to the rest of the Convention Area	Altered structure of future stock assessments	If 4B is found to be functionally isolated, a separate assessment may be constructed for that IPHC Regulatory Area	Genetics and	Population structure
Biological input	Assignment of individuals to source populations and assessment of distribution changes	urce populations and sessment of distribution lmprove estimates by Biological Region Will be used to define management targets for minimum spawning biomass by Biological Region		Genomics	Distribution
	Improved understanding of larval and juvenile distribution	or productivity	Will be used to generate potential recruitment covariates and to inform minimum spawning biomass targets by Biological Region	Migration	Larval and juvenile connectivity studies
Assessment data collection	Sex ratio-at-age	Scale biomass and	Annual sex-ratio at age for the commercial fishery fit by the stock assessment		Sex ratio of current commercial landings
and processing	Historical sex ratio-at-age	fishing intensity	Annual sex-ratio at age for the commercial fishery fit by the stock assessment	Reproduction	Historical sex ratios based on archived otolith DNA analyses
2. Assessment data collection and processing		Improve mortality accounting	May reduce depredation mortality, thereby increasing available yield for directed fisheries. May also be included as another explicit source of mortality in the stock assessment and mortality limit setting process depending on the estimated magnitude	Mortality and survival assessment	Whale depredation accounting and tools for avoidance
1. Fishery yield	Physiological and behavioral responses to fishing gear	Reduce incidental mortality	May increase yield available to directed fisheries	Mortality and survival assessment	Biological interactions with fishing gear
2. Fishery yield		Improve estimates of unobserved mortality	May reduce discard mortality, thereby increasing available yield for directed fisheries	Mortality and survival assessment	Best handling practices: recreational fishery

C. List of ranked biological uncertainties and parameters for management strategy evaluation (MSE) and their links to research areas and activities contemplated in the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21)

MSE Rank	Research outcomes	Relevance for MSE	Research Area	Research activities
Biological parameterization and	Improved understanding of larval and juvenile distribution	Improve parametization of the	Migration	Larval and juvenile connectivity studies
validation of movement estimates	Stock structure of IPHC Regulatory Area 4B relative to the rest of the Convention Area	Operating Model		Population structure
Biological parameterization and	Assignment of individuals to source populations and assessment of distribution changes	Improve simulation of recruitment variability and parametization of recruitment distribution in the Operating Model	Genetics and Genomics	Distribution
validation of recruitment variability and distribution	Establishment of temporal and spatial maturity and spawning patterns	Improve simulation of recruitment variability and parametization of recruitment distribution in the Operating Model	Reproduction	Recruitment strength and variability
3. Biological	Identification and application of markers for growth pattern evaluation			
parameterization and validation for growth	Environmental influences on growth patterns	Improve simulation of variability and allow for scenarios investigating climate change	Growth	Evaluation of somatic growth variation as a driver for changes in size-at-age
projections	Dietary influences on growth patterns and physiological condition			
Fishery parameterization	Experimentally-derived DMRs	Improve estimates of stock productivity	Mortality and survival assessment	Discard mortality rate estimate: recreational fishery



D. External funding received during the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21):

Project #	Grant agency	Project name	PI	Partners	IPHC Budget (\$US)	Management implications	Grant period
1	Saltonstall-Kennedy NOAA	Improving discard mortality rate estimates in the Pacific halibut by integrating handling practices, physiological condition and post-release survival (NOAA Award No. NA17NMF4270240)	IPHC	Alaska Pacific University	\$286,121	Bycatch estimates	September 2017 – August 2020
2	North Pacific Research Board	Somatic growth processes in the Pacific halibut (Hippoglossus stenolepis) and their response to temperature, density and stress manipulation effects (NPRB Award No. 1704)	ІРНС	AFSC-NOAA-Newport, OR	\$131,891	Changes in biomass/size-at-age	September 2017 – February 2020
3	Bycatch Reduction Engineering Program - NOAA	Adapting Towed Array Hydrophones to Support Information Sharing Networks to Reduce Interactions Between Sperm Whales and Longline Gear in Alaska	Alaska Longline Fishing Association	IPHC, University of Alaska Southeast, AFSC- NOAA	-	Whale Depredation	September 2018 – August 2019
4	Bycatch Reduction Engineering Program - NOAA	Use of LEDs to reduce Pacific halibut catches before trawl entrainment	Pacific States Marine Fisheries Commission	IPHC, NMFS	-	Bycatch reduction	September 2018 – August 2019
5	National Fish & Wildlife Foundation	Improving the characterization of discard mortality of Pacific halibut in the recreational fisheries (NFWF Award No. 61484)	ІРНС	Alaska Pacific University, U of A Fairbanks, charter industry	\$98,902	Bycatch estimates	April 2019 – November 2021
6	North Pacific Research Board	Pacific halibut discard mortality rates (NPRB Award No. 2009)	IPHC	Alaska Pacific University,	\$210,502	Bycatch estimates	January 2021 –March 2022
7	Bycatch Reduction Engineering Program - NOAA	Gear-based approaches to catch protection as a means for minimizing whale depredation in longline fisheries (NA21NMF4720534)	ІРНС	Deep Sea Fishermen's Union, Alaska Fisheries Science Center-NOAA, industry representatives	\$99,700	Mortality estimations due to whale depredation	November 2021 – October 2022
8	North Pacific Research Board	Pacific halibut population genomics (NPRB Award No. 2110)	ІРНС	Alaska Fisheries Science Center-NOAA	\$193,685	Stock structure	December 2021- January 2024
				Total awarded (\$)	\$1,020,801		

E. Publications in the peer-reviewed literature resulting from the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21):

2020:

Fish, T., Wolf, N., Harris, B.P., Planas, J.V. A comprehensive description of oocyte developmental stages in Pacific halibut, *Hippoglossus stenolepis*. *Journal of Fish Biology*. 2020. 97: 1880-1885. https://doi.org/10.1111/jfb.14551.

2021:


- Carpi, P., Loher, T., Sadorus, L., Forsberg, J., Webster, R., Planas, J.V., Jasonowicz, A., Stewart, I. J., Hicks, A. C. Ontogenetic and spawning migration of Pacific halibut: a review. *Rev Fish Biol Fisheries*. 2021. https://doi.org/10.1007/s11160-021-09672-w.
- Kroska, A.C., Wolf, N., Planas, J.V., Baker, M.R., Smeltz, T.S., Harris, B.P. Controlled experiments to explore the use of a multi-tissue approach to characterizing stress in wild-caught Pacific halibut (*Hippoglossus stenolepis*). *Conservation Physiology* 2021. 9(1):coab001. https://doi:10.1093/conphys/coab001.
- Loher, T., Bath, G. E., Wischniowsky, S. The potential utility of otolith microchemistry as an indicator of nursery origins in Pacific halibut (*Hippoglossus stenolepis*) in the eastern Pacific: the importance of scale and geographic trending. *Fisheries Research*. 2021. 243: 106072. https://doi.org/10.1016/j.fishres.2021.106072.
- Lomeli, M.J.M., Wakefield, W.W., Herrmann, B., Dykstra, C.L., Simeon, A., Rudy, D.M., Planas, J.V. Use of Artificial Illumination to Reduce Pacific Halibut Bycatch in a U.S. West Coast Groundfish Bottom Trawl. *Fisheries Research*. 2021. 233: 105737. doi: 10.1016/j.fishres.2020.105737.
- Sadorus, L., Goldstein, E., Webster, R., Stockhausen, W., Planas, J.V., Duffy-Anderson, J. Multiple life-stage connectivity of Pacific halibut (*Hippoglossus stenolepis*) across the Bering Sea and Gulf of Alaska. *Fisheries Oceanography*. 2021. 30:174-193. doi: https://doi.org/10.1111/fog.12512.

2022:

- Fish, T., Wolf, N., Smeltz, T. S., Harris, B. P., and Planas, J. V. Reproductive Biology of Female Pacific Halibut (*Hippoglossus stenolepis*) in the Gulf of Alaska. *Frontiers in Marine Science* 2022. 9:801759. doi: 10.3389/fmars.2022.801759.
- Jasonowicz, A.C., Simeon, A., Zahm, M., Cabau, C., Klopp, C., Roques, C., Iampietro, C., Lluch, J., Donnadieu, C., Parrinello, H., Drinan, D.P., Hauser, L., Guiguen, Y., Planas, J.V. Generation of a chromosome-level genome assembly for Pacific halibut (*Hippoglossus stenolepis*) and characterization of its sex-determining genomic region. *Molecular Ecology Resources*. 2022. In Press. doi: https://doi.org/10.1111/1755-0998.13641.
- Loher, T., Dykstra, C.L., Hicks, A., Stewart, I.J., Wolf, N., Harris, B.P., Planas, J.V. Estimation of postrelease longline mortality in Pacific halibut using acceleration-logging tags. *North American Journal of Fisheries Management*. 2022. 42: 37-49. DOI: http://dx.doi.org/10.1002/nafm.10711.




- F. Flow chart of progress resulting from the IPHC 5-Year Biological and Ecosystem Science Research Plan (2017-21) by research area leading to the IPHC 5-Year Program of Integrated Research and Monitoring (2022-2026)
 - 1. Migration and Distribution



Research outcomes: • Post-settlement migration from BS to GOA


2. Reproduction



Staff involved: Teresa Fish, MSc APU (2018-2020), Crystal Simchick, Ian Stewart, Allan Hicks, Josep Planas Funding: IPHC (2018-2020)

Publications (2): Fish et al. (2020) J. Fish Biol. 97: 1880–1885; Fish et al. (2022) Front. Mar. Sci. 9:801759

3. Growth



Research outcomes: • Effects of temperature on growth rates


• Temperature-specific molecular responses

Staff involved: Andy Jasonowicz, Crystal Simchick, Josep Planas

Funding: NPRB Grant#1704 (Sept. 2017-Feb. 2020)

Publications: Planas et al. (in preparation)

4. Mortality and Survival Assessment




Staff involved: Claude Dykstra, Allan Hicks, Ian Stewart, Josep Planas

Funding (3): Saltonstall-Kennedy NOAA (Sept. 2017-Aug. 2020); NFWF (Apr. 2019-Nov. 2021); NPRB#2009 (Jan. 2021-Mar. 2022)

Publications (2): Kroska et al. (2021) Conserv. Physiol.; Loher et al. (2022) North Amer. J. Fish. Manag. 42: 37-49

5. Genetics and Genomics



SNP detection and genotyping

F

Research outcomes:

- Sequenced genome (size=602 Mbp)
- Full annotation (NCBI) (27,944 genes)
- 24 chromosome-length scaffolds

Staff involved: Andy Jasonowicz, Josep Planas

Funding: IPHC, NPRB#2110

Publications: Jasonowicz et al. (2022) Mol. Ecol. Resour. (In Review)

APPENDIX II

Biological research areas in the 5-Year Program of Integrated Research and Monitoring (2022-2026) and ranked relevance for stock assessment and management strategy evaluation (MSE)

Research areas	Research activities	Research outcomes	Relevance for stock assessment	Relevance for MSE	Specific analysis input	SA Rank	MSE Rank	Research priorization
	Population structure	Population structure in the Convention Area	Altered structure of future stock assessments		If 4B is found to be functionally isolated, a separate assessment may be constructed for that IPHC Regulatory Area	Biological input	Biological parameterization and	2
Migration and population dynamics	Distribution	Assignment of individuals to source populations and assessment of distribution changes	Improve estimates of productivity	Improve parametization of the Operating Model	Will be used to define management targets for minimum spawning biomass by Biological Region	3. Biological input	validation of movement estimates and recruitment distribution	2
	Larval and juvenile connectivity studies	Improved understanding of larval and juvenile distribution	Improve estimates of productivity		Will be used to generate potential recruitment covariates and to inform minimum spawning biomass targets by Biological Region	3. Biological input	Biological parameterization and validation of movement estimates	2
	Histological maturity assessment	Updated maturity schedule			Will be included in the stock assessment, replacing the current schedule last updated in 2006			1
	Examination of potential skip spawning	Incidence of skip spawning	Scale biomass and	Improve simulation of	Will be used to adjust the asymptote of the maturity schedule, if/when a time- series is available this will be used as a direct input to the stock assessment	1. Biological		1
Reproduction	Fecundity assessment	Fecundity-at-age and -size information	reference point estimates	spawning biomass in the Operating Model	Will be used to move from spawning biomass to egg-output as the metric of reproductive capability in the stock assessment and management reference points	input		1
	Examination of accuracy of current field macroscopic maturity classification	Revised field maturity classification			Revised time-series of historical (and future) maturity for input to the stock assessment			1
		Identification and application of markers for growth pattern evaluation			May inform yield-per-recruit and other spatial evaluations of productivity that support mortality limit-setting			5
Growth	Evaluation of somatic growth variation as a driver for changes in size-at-age	Environmental influences on growth patterns	Scale stock productivity and reference point estimates	Improve simulation of variability and allow for scenarios investigating climate change	May provide covariates for projecting short-term size-at-age. May help to delineate between effects due to fishing and those due to environment, thereby informing appropriate management response		Biological parameterization and validation for growth projections	5
		Dietary influences on growth patterns and physiological condition			May provide covariates for projecting short-term size-at-age. May help to deleineate between effects due to fishing and those due to environment, thereby informing appropriate management response			5
	Discard mortality rate estimate: longline fishery	Experimentally-derived			Will improve estimates of discard mortality, reducing potential bias in stock assessment results and management of mortality limits	4 Fishers deld		4
Mortality and survival assessment	Discard mortality rate estimate: recreational fishery	DMR	Improve trends in unobserved mortality	Improve estimates of stock productivity	Will improve estimates of discard mortality, reducing potential bias in stock assessment results and management of mortality limits	Fishery yield	Fishery parameterization	4
	Best handling and release practices	Guidelines for reducing discard mortality			May reduce discard mortality, thereby increasing available yield for directed fisheries	2. Fishery yield		4
Fishing technology	Whale depredation accounting and tools for avoidance	New tools for fishery avoidance/deterence; improved estimation of depredation mortality	Improve mortality accounting	Improve estimates of stock productivity	May reduce depredation mortality, thereby increasing available yield for directed fisheries. May also be included as another explicit source of mortality in the stock assessment and mortality limit setting process depending on the estimated magnitude	Assessment data collection and processing		3

APPENDIX III

List of ranked research priorities for stock assessment

SA Rank	Research outcomes	Relevance for stock assessment	Specific analysis input	Research Area	Research activities
	Updated maturity schedule		Will be included in the stock assessment, replacing the current schedule last updated in 2006		Histological maturity assessment
1. Biological	Incidence of skip spawning	Scale biomass and	Will be used to adjust the asymptote of the maturity schedule, if/when a time-series is available this will be used as a direct input to the stock assessment		Examination of potential skip spawning
input	Fecundity-at-age and -size information	reference point estimates	Will be used to move from spawning biomass to egg-output as the metric of reproductive capability in the stock assessment and management reference points	Reproduction	Fecundity assessment
	Revised field maturity classification		Revised time-series of historical (and future) maturity for input to the stock assessment		Examination of accuracy of current field macroscopic maturity classification
2. Biological input	Stock structure of IPHC Regulatory Area 4B relative to the rest of the Convention Area	Altered structure of future stock assessments	If 4B is found to be functionally isolated, a separate assessment may be constructed for that IPHC Regulatory Area		Population structure
3. Biological	Assignment of individuals to source populations and assessment of distribution changes	Improve estimates	Will be used to define management targets for minimum spawning biomass by Biological Region	Migration and population dynamics	Distribution
input	Improved understanding of larval and juvenile distribution	of productivity	Will be used to generate potential recruitment covariates and to inform minimum spawning biomass targets by Biological Region		Larval and juvenile connectivity studies
1. Assessment	Sex ratio-at-age	Scale biomass and	Annual sex-ratio at age for the commercial fishery fit by the stock assessment	Depreduction	Sex ratio of current commercial landings
data collection and processing	Historical sex ratio-at-age	fishing intensity	Annual sex-ratio at age for the commercial fishery fit by the stock assessment	Reproduction	Historical sex ratios based on archived otolith DNA analyses
2. Assessment data collection and processing	New tools for fishery avoidance/deterence; improved estimation of depredation mortality	Improve mortality accounting	May reduce depredation mortality, thereby increasing available yield for directed fisheries. May also be included as another explicit source of mortality in the stock assessment and mortality limit setting process depending on the estimated magnitude	Fishing technology	Whale depredation accounting and tools for avoidance
1. Fishery yield	Physiological and behavioral responses to fishing gear	Reduce incidental mortality	May increase yield available to directed fisheries	Fishing technology	Biological interactions with fishing gear
2. Fishery yield	Guidelines for reducing discard mortality	Improve estimates of unobserved mortality	May reduce discard mortality, thereby increasing available yield for directed fisheries	Mortality and survival assessment	Best handling practices: recreational fishery

APPENDIX IV

List of ranked research priorities for management strategy evaluation (MSE)

MSE Rank	Research outcomes	Relevance for MSE	Research Area	Research activities
Biological parameterization and	Improved understanding of larval and juvenile distribution	Improve parametization of the		Larval and juvenile connectivity studies
validation of movement estimates	Stock structure of IPHC Regulatory Area 4B relative to the rest of the Convention Area	Operating Model	Migration and population	Population structure
Biological parameterization and	Convention Area Convention Area Improve simulation of recruitment variability and parametization of recruitment distribution changes Improve simulation of recruitment variability and parametization of recruitment distribution in the Operating Model Improve simulation of		dynamics	Distribution
validation of recruitment variability and distribution	Establishment of temporal and spatial maturity and spawning patterns	Improve simulation of recruitment variability and parametization of recruitment distribution in the Operating Model	Reproduction	Recruitment strength and variability
3. Biological	Identification and application of markers for growth pattern evaluation			
parameterization and validation for growth	Environmental influences on growth patterns	Improve simulation of variability and allow for scenarios investigating climate change	Growth	Evaluation of somatic growth variation as a driver for changes in size-at-age
projections	Dietary influences on growth patterns and physiological condition	22 digating similate sharige		
Fishery parameterization	Experimentally-derived DMRs	Improve estimates of stock productivity	Mortality and survival assessment	Discard mortality rate estimate: recreational fishery

APPENDIX V

List of ongoing and planned research projects (Will be linked to the website)

Research Project #	Project Title	Abstract	Objectives	Deliverables	Progress report	5YPRIM Research area	Management implications	Specific inputs into management	Period of Performance	PI	Funding source	Budget	Research prioritization for SAMSE
1	Leveraging multiple genomic approaches to investigate population structure and dynamics of Pacific halibut	The Pacific halbut (Hipcoglossus sterolesis) is a key further species in the North Pacific Ocean cooystem that support important commercial, recreational and subsistence fisheries and that is managed as a single stock by the International Pacific Halbut Commission. The overarching goal of the present study is to advance our understanding of Pacific Halbut population structure and dynamics in a changing climate through the use of genomic approaches to inform fishery management. In particular, we seek to Internove our current understanding of stock situative among sewinging groups of Pacific halbut in the northeast Pacific Ocean by conducting low coverage whole genome resequencing, a method that allows the characterization of genomic variation at the highest resolution possible and with which we will establish a baseline of Pacific halbut genetic devisery. Subsequently, we will leverage the characterization of genomic will have been supported to the pacific halbut of defined genetic baseline distincts. The results will be a supported to the pacific halbut of the pacific halbut of the pacific baseline of subsequently reformation on stock composition in the Pacific halbut fishery, as well as provide a tool to monitor changes in distribution associated with formatic change.	structure in the northeast Pacific Ocean using low	If Establishment of a baseline of Posich halbut penetic diversity. The genomic data produced wil represent a detailed baseline of Posich halbut genetic structure and dhershy at neutral and adaptive markers over a large geographical scale (Oard of Aleaka, Aleahan Isands and Berland and over a broad temporal scale (Best 30 years). 2.Delineation of fire-scale Pacific halbut attock structure. 3. Assignment of individuals to source populations and assessment of distribution charges.	09/NPRB Interim Report July 2023/IPHC-2023- WM2023-12	Migration and Population Dynamics	Altered structure of future stock assessments and MSE operating models. 2. Improve estimates of productivity. 3. Improve understanding of population distribution and the effects of distributing fishing effort.	separate assessment may be constructed for that IPHC Regulatory Area. Research	12/01/2021- 02/16/2024	Josep Planas	External (North Pacific Research Board; Project No. 2110)	\$193,685	Priority Rank #2
2	habitat	The FHC Secretarist recently completed a study to investigate the connectivity between spawning grounds and possible settlement earse based on a biophysical lamel transport model (Sadorus et al., 2021; https://doi.org/10.1111/6g.12512), Although it is known that Pacific halbut, following the playing lamel phase begin here demonstrating as roughly formorth-old juveniles, settling in shallow nursery (settlement) areas, near or outside the mouths of bays (Cupil et al., 2021; https://doi.org/10.1071/1110-021-03072-030-0, way tills in ordination is available on the geographic formorth or the settlement of the settlement of the settlement areas for juvenile Pacific halbut throughout PHC Correction Waters.	presence. 2. Create a map of suitable settlement habitat by combining available bethymetry information (e.g. benthic sediment composition and shoreline morphological data) and information on recorded presence of age-0, age-1 and age-2 Pacific halibut juveniles as well as absence of young Pacific halibut noted by various nursery habitat projects focused on other fatfish species.	Map of juvenile Pacific halibut habitat.	IPHC-2023-SRB022- 09/IPHC-2023-WM2023- 12	Migration and Population Dynamics	improve estimates of productivity	Will be used to generate potential recruiment covariates and to inform minimum spawning biomass targets by Biological Region	01/01/2023- 12/31/2025	Josep Planas	Internal	\$0	Priority Rank #2
3	Female reproductive assessment	In fisheries, understanding the reproductive biology of a species is important for estimating the reproductive potential and spamning biomas of the stock and, consequently, for optimizing management of the species. Recent sensitivity analyses have shown the importance of changes in spamning output in female Pacific heliality due to changes in maintarty schedules; in feural pacific heliality of the pacific heliality, research efforts are being conducted to characteristic female reproductive biology of female Pacific hallout, research efforts are being conducted to characteristic female reproductive epacify in this species. Improved browdege again relation to the pacific hallout, research efforts are being conducted to characteristic female reproductive epacify in this species. Improved browdege gain relations to the reproductive characteristic productive characteristic productive characteristic productive capacity and success in this important species.	parameters that will be used to revise current estimates of female and male age-at-maturity. 3.Compare macroscopic (based on field	Lipdated maturity schedule coastwide. Zenoundlys-dag and -size estimates. Revised field maturity classification. Antermation on skip-spawring.	PHC-2023-SRB002- 00)PHC-2023-WM0023- 12	Reproduction	Scale biomass and reference print estimates, Improve estimates of spawning biomass in the stonck-was estimated and include assessment and invincion assessment and invincion biomass in the MSE operating model.	Research outcomes will be included in the stock assessment, replacing the current maturity schedule last updated in 2006, schedule last updated in 2006, schedule last updated in 2006, and a separate of the grant of the control of t	01/01/2017- 12/31/2026	Josep Planas	Internal	\$51,834 (FY2024)	Priority Rank #1
4	protection as a means for minimizing whatel depredation in longline fisheries	In the north Paulife, both foller (Crimius oras) and Sperm (Physeler macrocephalus) wheles are involved in deprediction behavior in Paulife inballed (Hippoglosus sendence) in 2011 and 2012 fisheries observers estimated that 6.9% of Paulife habitat sets were affected by whale depredation in the Berring Sare, Reductions in catch per unit effort (CPUE) when whalese were present ranged across ageographic regions from 515-57% for Paulife habitat. These impacts also incur significant fine, leut, and presented costs to failing operations. From a faitheries management perspective, depredation creates an additional and highly uncertain source of mortality, loss of date (e.g. compromised survey activity), and relocated fishery efficients accurate of mortality, loss of date (e.g. compromised survey activity), and relocated fishery efficients ourselved of mortality, loss of date (e.g. compromised survey activity), and relocated instead of the compromised survey activity, and relocated instead of the compromised survey activity, and relocated instead of the compromised survey activity, and relocated of the fisher of the compromised survey activity. As a survey and the compromised survey activity, and relocated to the fisher of the fisher of whale deprediction or activity rates. In the satisfies has seen survey, the same survey and the satisfies of the whales include physical injury date to being near vessels and gear. Gamptotion of social structure, and developing an artificial relations on food dams that can be affected by fishery dynamics. Meny efforts have been made over the constructed, deployed, or emanded without applicantly disrupting room in fairing operations, or disruptive activity of success and ease of adoption in each fishery bit more have provided a long-term solution. These are increasing data sources supporting the notion that behindogles which motion intal contract between pair and depredations will reduce the likelihood of fronging attempts around mortality and gear demange. Recent studies using physical c	designs that can be deployed effectively using current tongline fishing techniques.	protection devices in field tests and provide direction for them modification and larger scale experimental testing.			improve mortality accounting, improve estimates of stock productivity.	Research outcomes may reduce depredation mortality, thereby increasing available yield for directed fisheries. May also be included as another explicit source of mortality in the stock assessment and mortality limit settling process depending on the estimated magnitude.	10/30/2023	Stewart	External (Byeatch Reduction Engineering Program NOAA: Project NOA2 1NMF4720534)	\$99,700	Phordy Rank #3
5	Use of artificial intelligence (Al) for determining the age of Pacific haibut from images of collected toliths	The EMS Secretariat is boding at options for supplimenting current Peofic hallful ageing protos with automatical ageing that does not require extensive bodins research supplies and provided that the provided and the second supplies and the second supplies and the second supplies and the second supplies and supplies	aged otoliths 2.Train and validate a CNN model for automated ageing	Predictive CNN model for ageing Pacific halibut complementing traditional methods 2.4 report complementing traditional methods to traditional ageing techniques	N∕A	Age composition data (both fishery- dependent and fishery-independent)		Al-driven age determination offers a critical enhancement to stock assessment methodologies, ciking in the estimation of growth rates, maturity, and population structure of Pacific hallbut.	12/2024+	Barbara Hutniczak	Internal	\$0	Priority Rank #1



roposed			I	I									
Research Project #	Project Title	Abstract	Objectives	Deliverables	Progress report	5YPRIM Research area	Management implications	Specific inputs into management	Requested period of performance	PI	Targeted funding source	Requested budget	Research prioritization for SA/MSE
1	to inform population structure and dynamics affecting coastal communities	effort has so far resulted in the identification of 1.15, million autosomal single nucleotide oppomychnighmic (PSPs), of which if almois NSPs have a minor alled frequency higher than 0.05. Considerable progress is currently being made towards using genome approaches to setablish a generic baseline for the available systeming groups, and towards the development of genomic towards and the second of the second of the progress of the second of genetic samples from spawing groups of the WA coast limits the application of the above mentioned genomic tools to advance our understanding of population structure, movement, connectivity, adaptive characteristics, and environmental responses of Pacific habble in Convention waters. Although no major spawing ground has been mapped south of Cape St. James in the southern sign of Haad Cave (ICF Paris, 1963), archeological records along with what is now PHC Regulatory was A (Saliema-Harley, 2018). Additionally, corresponding of spawing pacific haibut in the winter of the WA coast indicate that Pacific haibut, at least historically, approach of spawing pacific haibut south of Cape Flattery and the existence of suitable spawing habilat for Pacific haibut with waters and they brought pacification and sprought pacification of the presence of spawing grounds for Pacific haibut of the WA coast. Therefore, the sicentification of potential wither spawing grounds for Pacific haibut of the WA coast. Therefore, the sicentification of potential wither spawing grounds for Pacific haibut of the WA coast. Therefore, the sicentification of potential wither spawing grounds for Pacific haibut of the WA coast the proposal is to characterize the generic composition of Pacific haibut of the WA coast using proposal is to characterize the generic composition of Pacific haibut and the WA coast using proposal is to characterize the generic composition of Pacific haibut and the WA coast using proposal is to	ecological knowledge and collect biological samples. Z. To characterize the reproductive condition of female and make Pacific hallout of the WA coast during the winter spawning season. 3. To generate and incorporate genomic data from winter spawning group of the WA coast to existing winter spawning group of the WA coast to existing seaso, in the north-eastern Pacific Ocean to establish areas in the north-eastern Pacific Ocean to establish areas in the north-eastern Pacific Ocean to establish an expanded baseline of Pacific halbut genetic diversity.	off the WA coast is californiation, pawmig or man collection of brobigoid (general and reproductive) samples. 2. Extended bearing of Practic halbut genetic and reproductive) samples. 2. Extended bearing of Practic halbut genetic ability of the scalar fractic halbut stock function in WA wasters and coast-wide.		Mgration and Population Dynamics	Alered structure of future stock assessment and MSE stock assessment and MSE operating models, improved estimates of productivity case twide.	Convention waters will inform management actions by validating management units. Research outcomes will be used to define management targets for minimum speaming biomass by Biological Region.	02/01/2024-1/31/2026	Josep Planas	External (Washington Sea Grant). Full proposal submitted in the May 2023. Proposal not existed for funding.		Priority Rank #2
2	Full scale testing of devices to minimize what depredation in longline flaheries	International Conference of the Conference of th	Assess the performance metrics of catch protection devices on the size, number, and condition of fish successfully entrained in the devices	Further define and develop previously detailled, pipel pricity work that can break the reward cycle of depredation behavior and thereby suppress is providence. 2 Bull don't strategies to protect the providence of the provid		Fishing technology	Improved accuracy of mortally estimates, improve estimates of productivity	Will be used to generate potential recruitment covariates and to inform minimum spawning biomass targets by Biological Region	11/1/2023-04/30/2025	Claude Dykstra/lan Stewart	External (Bycatch Reduction Engineering Program NOAA): Full proposal submitted in March 2023: Avanded.	\$199,870	Priority Rank #3
3	Development of a non-lethal genetic-based method for aging Pacific halibut	Recordation accusion with this case has been anosted (Posthed et al. 2021). Association that Richael Medical Commission (Posthed Medical Commission (Posthed Medical Commission) (Posthed C	halibut fin fissue. 2. To develop an age prediction model based on DNA methylation patterns: an epigenetic clock for Pacific halibut. 3. To develop a targeted DNA methylation assay for larger scale age estimations.	Reduced representation genome-wide map of DNA methylation a traingle bases-pair reaching Page 18 per periodicing model for Pacific halibut fin tissue. 2. App predicting model for Pacific halibut using fin tissue.		Mygration and Population Dynamics/Female Reproductive Assessment/Growth	Age is a critical input for stock assessment.	Age is a key blotogical input into stock assessment as it is used for estimating fish growth, fish maturity and fecundity-at-eag, and mortality raties as well as positions are successful as the successful and the successful as t	02/01/2024 1/31/2026	Josep Planas	External (Aleska See Grant), Full process submitted in Net 2023. Declarion expected January 2024.	\$60,374	Priority Rank #1

APPENDIX VI

Proposed schedule of outputs

	20	22	20	23	20	24	20	25	20	26
Biology and Ecology	•								•	
Migration and population dynamics										
Reproduction										
Growth										
Mortality and survival assessment										
Fishing technology										
Stock Assessment										
Management Strategy Evaluation										
Monitoring										

APPENDIX VII

Proposed schedule of funding and staffing indicators: Biology and Ecology

Research areas	Research activities	Required FTEs/Year	IPHC FTEs/Year		20	22	2023				20	2024				2025			20	26	IPHC Funds	Grant Funds	
	Larval and juvenile connectivity and early life history studies	0.45	0.45				R	B1			RB2											Yes	NPRB #2100
	Population structure	0.4	0.8				RB1															No	NPRB #2110
	Adult migration and distribution	0.4	0.0																			No	NPRB #2110
Migration and Population Dynamics	Close-kin mark-recapture studies	1	0																			No	Planned
	Seascape genomics	1	0																			No	Planned
	Genome-wide association analyses	1	0																			No	Planned
	Genomic-based aging methods	1	1					RS	1													Yes	No
	Maturity-at-age estimations	0.75	0																			Yes	No
	Fecundity assessment	0.5	0.25								RB4					I	RS 2					Yes	No
Reproduction	Examination of accuracy of current field macroscopic maturity classification	0.25	0.20																			Yes	No
	Sex ratio of current commercial landings	0.5	0.75	LT																		Yes	No
	Recruitment strength and variability	0.5	0													R	S 2					Yes	Planned
Growth	Environmental influences on growth patterns	0.5	0.5							MS	stu	dent										No	Planned
Clowin	Dietary influences on growth patterns and physiological condition	0.5	0.2							RB3												No	Planned
	Discard mortality rate estimate: recreational fishery	0.5																				No	NPRB #2009
Mortality and survival	Best handling practices: recreational fishery	0.5	1			RB 3																No	NPRB #2009
assessment	Whale depredation accounting and tools for avoidance	0.5	,																			No	BREP
IPHC staff (Planned):	Biological interactions with fishing gear	0.5																				No	BREP

IPHC staff (Planned):

- RS1: Research Scientist 1(PhD; Life History Modeler I). Full time temporary position (100% research;
- RS2: Research Scientist 1(PhD; Life History Modeler II). Full time temporary position (100% research;
- RB1: Research Biologist 1 (Geneticist; MSc). Full time temporary position (until April 2022; 1 FTE). 55% of salary covered by Grant NPRB#2110.
- RB2: Research Biologist 2 (Early Life History; MSc). Full time permanent position (40% research; 0.4 FTE)
- RB3: Research Biologist 3 (DMR; MSc). Full time permanent position (100% research; 1 FTE)
- RB4: Research Biologist 4 (Maturity and Fecundity; MSc). Full time permanent position (100% research; 1 FTE)
- LT: Laboratory Technician (MSc). Full time temporary position (100% research; 1 FTE)